

Université de Sherbrooke

**Photodynamic Therapy: Enhancement of phthalocyanine targeting
from modifications on the macrocycle to the use of protein
delivery vehicles**

By

Cynthia M. Allen

Department of Nuclear Medicine and Radiobiology,
Faculty of Medicine, University of Sherbrooke,
Sherbrooke, Québec, Canada, J1H 5N4

Thèse présentée à la Faculté de Médecine
en vue de l'obtention du grade de
Philosophiae Doctor (Ph.D.) en radiobiologie

June 11, 2001



National Library
of Canada

Acquisitions and
Bibliographic Services

395 Wellington Street
Ottawa ON K1A 0N4
Canada

Bibliothèque nationale
du Canada

Acquisitions et
services bibliographiques

395, rue Wellington
Ottawa ON K1A 0N4
Canada

Your file Votre référence

Our file Notre référence

The author has granted a non-exclusive licence allowing the National Library of Canada to reproduce, loan, distribute or sell copies of this thesis in microform, paper or electronic formats.

The author retains ownership of the copyright in this thesis. Neither the thesis nor substantial extracts from it may be printed or otherwise reproduced without the author's permission.

L'auteur a accordé une licence non exclusive permettant à la Bibliothèque nationale du Canada de reproduire, prêter, distribuer ou vendre des copies de cette thèse sous la forme de microfiche/film, de reproduction sur papier ou sur format électronique.

L'auteur conserve la propriété du droit d'auteur qui protège cette thèse. Ni la thèse ni des extraits substantiels de celle-ci ne doivent être imprimés ou autrement reproduits sans son autorisation.

0-612-85168-0

Canada

*It is not what we know that is important,
it is what we do not know.*

Charles F. Kettering

*A task without a vision is drudgery.
A vision without a task is a dream.
A task with a vision is victory.*

To Wesley,

Table of Contents

List of illustrations

Figures.....	i
Tables.....	ii

List of Abbreviations and Symbols.....	iii
--	-----

Abstract	v
----------------	---

Chapter 1 - Introduction	1
1.1-Historical perspective.....	3
1.2-Mechanisms of photosensitization	4
1.3-First generation photosensitizers: Hematoporphyrin derivatives.....	6
1.4-Second generation photosensitizers: Phthalocyanines	9
1.4.1-Phthalocyanine induced apoptosis or necrosis	12
1.4.2-Phthalocyanine delivery systems.....	15
1.5-Research objectives.....	18

Chapter 2

Photodynamic properties of amphiphilic derivatives of aluminum tetrasulfo phthalocyanine	20
---	----

Chapter 3

Low density lipoprotein-bound aluminum sulfophthalocyanine: Targeting tumor cells for photodynamic therapy.....	30
--	----

Chapter 4

Photodynamic therapy: Tumor targeting with adenoviral proteins.....	38
---	----

Chapter 5

Attenuation of photodynamically induced apoptosis by an RGD containing peptide	51
---	----

Chapter 6 – Discussion	61
6.1-Amphiphilic phthalocyanines	62
6.2-Protein carriers.....	63
6.3-Serum Proteins.....	64
6.3.1-Albumin.....	65
6.3.2-Lipoproteins	66
6.4-Antibody conjugates.....	71
6.4.1-Phthalocyanines	71
6.5-Epidermal growth factor	74
6.6-Adenoviral carriers.....	75
6.7-Advances toward peptidic carriers.....	77
6.8-Conclusions	80
 Chapter 7 - Acknowledgements	 84
 Chapter 8 - Bibliography	 85
 Appendices	
I-Role of Activated Oxygen Species in Photodynamic Therapy	117
II-Photodynamic Therapeutics: Basic Principles and clinical applications	143
III-Current Status of Phthalocyanines in the Photodynamic Therapy of Cancer.....	155

List of Figures

1.1	Type I and Type II photosensitized oxidation reactions.....	6
2.1	AlPcS ₄ (C _n) structure and absorption spectrum.....	22
2.2	AlPcS ₄ (C8) absorption spectra in PBS, methanol, 1% CRM	24
2.3	A549 cell membrane adsorption of AlPcS ₄ derivatives.....	24
2.4	Effect of LDL on Pc uptake in A549 cells.....	25
2.5	A549 photocytotoxicity using AlPcS ₄ derivatives.....	25
3.1	Diagram depicting AlPcS ₄ coupling to LDL.....	32
3.2	Effect of light dose cell survival with LDL coupled AlPcS ₄	34
3.3	Effect of AlPcS ₄ -LDL concentration on cell survival	34
4.1	PAGE and Western analysis of Ad proteins coupled to a Pc.....	42
4.2	AlPcS ₄ A ₁ and AlPcS ₄ A ₂ structure.....	42
4.3	A549 cell adsorption of AlPcS ₄ coupled to adenoviral proteins	43
4.4	A549 photocytotoxicity using AlPcS ₄ -Ad2 proteins.....	44
5.1	Effect of RGD on AlPcS _{2adj} phototoxicity	54
5.2	Effect of temperature on RGD attenuation of apoptosis.....	55
5.3	Effect of rinsing cells post peptide prior to PDT.....	55
5.4	Effect of overnight incubation with RGD on cell survival post PDT	56
5.5	RGD effect on nucleosomal laddering induced by PDT and UV-C.....	57
5.6	RGD effect on PDT induced apoptotic bodies.....	57

List of Tables

2.1	L-Tryptophan photooxidation by AlPcS ₄ derivatives	23
2.2	Phototoxic activity of AlPcS ₄ derivatives versus retention times	25
2.3	EMT-6 tumor response <i>in vivo</i> using AlPcS ₄ derivatives.....	26
3.1	Photoinactivation of EMT-6 and A549 with LDL-Pc	35
3.2	EMT-6 tumor response post PDT with LDL-Pc	35
4.1	Labeling efficiency of viral proteins with AlPcS ₄	44
4.2	Phototoxic activity of AlPcS ₄ -Ad2 protein conjugates.....	44
4.3	EMT-6 tumor response using AlPcS ₄ -Ad2 protein conjugates.....	47
6.1	Summary of photosensitizer serum-based vehicles.....	70
6.2	Summary of photosensitizer-antibody conjugates.....	73

Abbreviations and Symbols

A549 cells	adenocarcinoma lung cancer cells
Ad2wt	wildtype adenovirus serotype 2
AlPcS	aluminum sulfophthalocyanine
AlPcS _{2adj}	aluminum disulfophthalocyanine
AlPcS ₄	aluminum tetrasulfophthalocyanine
AlPcS ₄ (C _n)	aluminum mono-(alkylaminosulfonyl)-tetrasulfophthalocyanine
AlPcS ₄ A ₁	aluminum mono(6-carboxypentylaminosulfonyl)tetrasulfophthalocyanine
AlPcS ₄ A ₂	aluminum di(6-carboxypentylaminosulfonyl)tetrasulfophthalocyanine
B _{max}	receptor density
ATP	adenosine triphosphate
BPD	benzoporphyrin derivative
BSA	bovine serum albumin
CRM	Cremophor EL
DDM	aspartate-aspartate-methionine peptidic sequence
DMEM	Dulbecco's modified Eagle's medium
DMF	N,N-dimethylformamide
ECM	extracellular matrix
EDTA	ethylenediamine tetraacetic acid
EGF	epidermal growth factor
EMT-6 cells	murine mammary tumor cells
ϵ	epsilon, molar extinction coefficient
ERK	extracellular regulating kinase
FAK	focal adhesion kinase
FCS	fetal calf serum
HSA	human serum albumin
HEP-2 cells	human epidermoid laryngeal carcinoma
HpD	hematoporphyrin derivative
HPPI	3 α -hydroperoxy-1,2,3,3 α ,8,8 α -hexahydropyrrolo[2,3 β]indole-2- carboxylic acid

IC	internal conversion
ISC	intersystem crossing
LDL	low-density lipoprotein
λ_{max}	maximum absorption wavelength
MOI	multiplicity of infection
MTT	3-(4,5-dimethylthiazol-2-yl)-2,5-(diphenyltetrazolium bromide)
O. D.	optical density
PAGE	polyacrylamide gel electrophoresis
PBS	phosphate-buffered saline
PDT	photodynamic therapy
PEG	polyethylene glycol
PFU	plaque forming unit
ϕ	quantum yield
PI-3 kinase	phosphatidylinositol 3-kinase
PLA	poly(D,L-lactic acid)
RGD	arginine-glycine-aspartic acid peptidic sequence
ROS	reactive oxygen species
R_t	retention time
S_0	ground state of sensitizer
S_1	singlet excited state of sensitizer
SDS	sodium dodecyl sulfate
$^1\text{O}_2$	singlet oxygen
Sens*	excited state of sensitizer
Sf9	<i>Spodoptera frugiperda</i>
τ_T	lifetime of triplet state of sensitizer
T_1	triplet excited state of sensitizer
TAE	Tris-acetate EDTA
TE	10 mM Tris pH 7.5, 1 mM EDTA
TFA	trifluoroacetic acid
TNF	tumor necrosis factor

Abstract- Photodynamic therapy is a promising treatment modality that involves the localization of a light-sensitive drug or photosensitizer in the target tissue prior to illumination with visible light. In the presence of molecular oxygen, cytotoxic agents are generated upon illumination and trigger a cascade of biochemical responses that may eradicate both malignant and non-malignant conditions. Phthalocyanine derivatives are second generation photosensitizers with enhanced photophysical, photochemical and photobiological properties as compared to the clinically accepted hematoporphyrin derivative Photofrin[®]. The current study aims at improving the targeting of these photosensitizers to cancerous tissues either through modifications on the phthalocyanine macrocycle or through protein carriers. AlPcS₄ derivatives substituted with long aliphatic chains of varying lengths were photodynamically active *in vitro* against a lung adenocarcinoma cell line, A549 cells. Activity varied directly with the degree of lipophilicity: AlPcS₄(C16) > AlPcS₄(C12) > AlPcS₄(C8) > AlPcS₄(C4). *In vivo* studies using the EMT-6 murine model were promising, requiring 0.2 $\mu\text{mol kg}^{-1}$ and 400 J cm⁻² to completely ablate the tumor. LDL receptor expression is often upregulated in cancer cell membranes. Therefore, in order to enhance Pc targeting, LDL was loaded with AlPcS by two different approaches. The AlPcS₄(C12) derivative was non-covalently inserted in the phospholipid bilayer of LDL particles. Conversely, AlPcS₄ substituted with two caproic acid spacer chains were covalently attached to the LDL protein moiety. While the latter had only marginal success, the former proved to be photodynamically active both *in vitro* and *in vivo*. To further exploit cell surface receptors as potential targets in PDT, adeno-viral proteins were covalently labeled with AlPcS₄. Adenoviruses penetrate the cell via receptor mediated endocytosis. The Ad penton base contains 5 repeats of the peptidic RGD binding sequence. This motif binds with high affinity to several members of the integrin receptor family. Integrin receptors play a major role in tumorigenesis and metastases with altered expression being a factor. The class of integrin receptors with high affinity for the RGD sequence is expressed in high numbers on A549 cells. Cell uptake of the AlPcS₄-adenovirus protein conjugates was enhanced over the parental Pc; however, photocytotoxicity results were unfavorable. *In vivo* studies using the same rodent model as that of the LDL study were encouraging. The AlPcS₄-adenovirus protein conjugates caused complete tumor regression at a dose of 0.5

$\mu\text{mol kg}^{-1}$ and 400 J cm^{-2} . The role of RGD on PDT induced cell death was also investigated. It was anticipated that soluble RGD would have a synergistic effect on PDT induced apoptosis yet the opposite was found. RGD enhanced cell survival following PDT in a dose dependent manner in both a receptor negative (EMT-6) and a receptor positive (A549) cell line. This was an unexpected phenomenon and warrants further investigation. Once understood, the possibility of designing peptidic carriers for photosensitizers or adjunct therapies using the RGD peptide are within the realm of possibility. Nonetheless, specific targeting of AlPcS₄ derivatives to cancer cells resulted in enhanced photodynamic activity as compared to the parental molecule, AlPcS₄.

Chapter 1.

Introduction

1. Introduction

Cancer is one of the leading causes of death and disease in the Western World. Over 1.2 million new cases of cancer were diagnosed in the United States in 1999 (American Cancer Society) while approximately one third of the population of Western Europe can expect to be touched by the disease within their lifetime (Brown et al., 1999). Cancer is an unregulated proliferation of transformed cells displaying an altered phenotype, distorted differentiation and genetic instability. These cells usually spread to and infiltrate distant sites of the body, giving rise to metastases (Fingert et al, 1997). Traditional cancer treatment modalities such as surgery, chemotherapy and radiation therapy aim at locoregional or systemic control of the malignancy and involve a delicate balance between removing or destroying diseased tissue while sparing neighboring healthy cells. These conventional therapies produce serious side effects caused by the loss of normal cell function as a result of their relatively indiscriminate cytotoxic effects. Furthermore, despite operating at near optimal efficiencies, these treatment protocols often are not sufficiently effective, frequently leading to important recurrences. While therapeutic improvements are continually being made, there have been no substantial enhancements in survival for a number of major malignancies over the past decade (Brown et al., 1999). Consequently, the search for new and improved treatment protocols that exhibit more selectivity for diseased tissue and less morbidity remains unabated.

Photodynamic therapy (PDT) is a promising new treatment modality that has recently been accepted in clinic as a curative and/or palliative therapy for cancer and also for non-malignant disorders. Photodynamic therapy entails a classic binary system which combines a light-activated drug (photosensitizer) with light of the appropriate wavelength

that is consistent with the absorption spectrum of the given photosensitizer. Illumination of the photosensitizer typically with visible light triggers a sequence of photophysical, photochemical and photobiological processes that cause irreversible photodamage to the target tissue. Neither the photosensitizer nor the light are harmful by themselves and only produce cytotoxic species when combined in the presence of molecular oxygen. Like all new cancer therapies, PDT primarily aims at destroying diseased tissue without causing inordinate injury to surrounding healthy cells. This targeting of diseased tissue is aided by a number of factors unique to PDT. The dual selectivity of PDT is produced by both a preferential uptake and/or prolonged retention of the photosensitizer by the diseased tissue along with the ability to confine activation of the photosensitizer to the target volume. This is accomplished by restricting the illumination to that specific volume, often using lasers and fiber optics. Hence, PDT allows for the selective destruction of tumors, providing the potential for minimal morbidity and large therapeutic indices.

1.1 Historical perspective

The therapeutic benefits of light have been known since the ancient world with early Greek, Egyptian, Roman and Indian civilizations all recognizing the healing powers of the sun. In India, as early as 1400 B.C., extracts of *Psoralea corylifolia* were given orally, followed by exposure to sunlight, in order to treat vitiligo (Daniell and Hill, 1991; Bonnett, 1999). Modern photodynamic therapy however can trace its origins to the beginning of the 20th century when a medical student, Oscar Raab, working under the direction of Dr. von Tappeiner in Munich, noticed the lethal effects of light on *Paramecia* treated with acridine dyes (Raab, 1900). This phenomenon was more extensively studied

by von Tappeiner and Jodlbauer who coined the term “photodynamic action” (von Tappeiner and Jodlbauer, 1907). It was noted that other compounds such as eosin were able to induce rapid cell kill in the presence of light. It was established that dissolved molecular oxygen is needed for the reaction to occur. The first attempt at using PDT as a cancer treatment dates to 1903 where topically applied eosin was used in conjunction with sunlight to treat skin cancer (von Tappeiner and Jesionek, 1903). The relative specificity of PDT was clearly demonstrated by Lipson in 1961 when the use of hematoporphyrin derivative (HpD) for the fluorescent detection of tumor tissue was reported (Lipson et al., 1961). In 1975, Dougherty first reported the eradication of transplanted animal tumors with HpD and red light without excessive damage to surrounding skin (Dougherty et al., 1975). This was soon followed by the first systemic clinical trials in humans and since then the field of PDT has rapidly expanded (Dougherty et al., 1978). Due to the clinical acceptance of Photofrin[®], verteporfin and the prodrug 5-aminolaevulinic acid (ALA), PDT has evolved into a useful treatment modality.

1.2 Mechanisms of photosensitization

Upon illumination, the photosensitizer is excited from its electronic ground state (S_0) to its first excited state (S_1). This excited singlet state has a lifetime in the nanosecond range (Phillips, 1997), which is far too short for effective interaction with surrounding biological molecules. The S_1 state can dissipate its excitation energy via radiative fluorescence or non-radiative internal conversion (IC). The most important of these processes in terms of PDT, however, is intersystem crossing (ISC) to the first excited triple state (T_1) with the corresponding change in electron spin. The lifetime of the T_1 state is typically in the μ s to ms range since the return $T_1 \rightarrow S_0$ is spin-forbidden

(Ochsner, 1997). The large increase in lifetime allows for the excited photosensitizer to interact with its environment and it is generally accepted that the triplet state is responsible for the generation of the cytotoxic species produced during PDT in the majority of cases (Sharman et al., 2000).

This excited triplet state can interact in one of two ways, defined as Type I and Type II (Figure 1). The T_1 state of the excited photosensitizer can be quenched via a hydrogen atom extraction or an electron transfer reaction with neighboring substrates. These reactions termed a Type I mechanism, lead to the formation of radicals and radical ions. These radical species react efficiently with molecular oxygen, leading to important oxidative damage and the production of reactive oxygen species such as superoxide anion, hydrogen peroxide and the hydroxyl radical. The following radical chain reactions cause important biological lesions. On the other hand, the T_1 state can transfer its energy to molecular oxygen via a triplet-triplet annihilation reaction, generating singlet oxygen (Sharman et al., 2000). This Type II reaction mechanism has been shown to be predominant during PDT. Singlet oxygen, the most important cytotoxic agent generated during PDT, is highly reactive due to its zwitterionic character and can readily react with a number of biologically important substrates including amino acids, phospholipids, and nucleic acids (Singh, 1982). This leads to disruption of cellular and mitochondrial membranes, increased cellular permeability, the loss of vital protein and enzyme function and irreparable DNA damage, all of which can be lethal to the cell. It is generally accepted that the Type II mechanism prevails during PDT and that singlet oxygen is the most important cytotoxic species produced (Sharman et al., 2000; Weishaupt et al., 1976; Valenzano, 1987; Fuchs and Thiele, 1998). The range of singlet oxygen in cellular

medium is limited to approximately 45 nm (Moan and Boye, 1981 and Moan, 1990). With the diameter of human cells ranging from 10^4 to 10^5 nm, the site of primary generation of singlet oxygen consequently determines which subcellular target is attacked, either initiating an apoptotic or necrotic response. Appendix I outlines the importance of singlet oxygen in photodynamic therapy. However, while it is known that Type II processes predominate during PDT, Type I mechanisms become important at low oxygen concentrations and in more polar environments (Ochsner, 1997; Sharman et al., 2000). These Type I mechanisms are no doubt involved in most photodynamic processes to a certain degree. Irrespective of the initial reaction, both Type I and Type II mechanisms lead to similar oxidative damage and comparable free radical chain reactions. The two processes, Type I and Type II are outlined in figure 1.




Figure 1. Diagrammatic presentation of Type I and Type II photosensitized oxidation reactions (Adapted from C. S. Foote, 1991)

1.3 First generation photosensitizer: Hematoporphyrin derivative

The first photosensitizers accepted for clinical use by governmental regulatory agencies are the first generation hematoporphyrin derivatives such as Photofrin®.

Photofrin[®], an enriched sample of the more active dimeric and oligomeric components, has been accepted in several countries for the treatment of early- and late-stage lung cancer, superficial and advanced oesophageal cancer, bladder cancer, superficial and early-stage gastric cancer, early stage cervical cancer and cervical dysplasia. In addition, Photofrin[®] is being investigated as a possible therapy for a number of other malignant and non-malignant conditions. Other hematoporphyrin derivatives are also being used, with Photoheme accepted for clinical use in Russia for easily accessible cancers including skin, breast, oropharyngeal, lung, larynx and gastrointestinal cancers along with psoriasis and recurrent blindness (Sharman et al., 1999; Sobolev and Stranadko, 1997; Stranadko et al., 1996; Sokolov et al., 1994).

Despite the clinical success achieved using Photofrin[®], this first generation photosensitizer has several important drawbacks. It is a complex chemical mixture produced by the reaction of hematoporphyrin with 5% sulfuric acid in acetic acid followed by treatment with aqueous base and neutralization (Bonnett, 1995). The resulting dimers and oligomers, linked primarily with ester and ether bonds, may vary with different preparations and storage times and make structure-activity relationships impossible to determine (Brown and Truscott, 1993). It absorbs very weakly at the therapeutic wavelength of 630 nm, where tissue penetration of light is minimal, limiting treatment to tumor depths of no more than 5 mm (Wöhrle et al., 1998). While suitable for more superficial lesions presently being treated, absorption at longer wavelengths is necessary to treat more deep-seated or larger tumors. In addition, hematoporphyrin derivatives have proven to be ineffective for such cancers as pigmented melanoma due to overlapping absorption spectra of the photosensitizer and melanin in the malignant tissue

(Peeva et al., 1999). Most importantly, hematoporphyrin derivatives exhibit an extended retention in cutaneous tissue for up to 10 weeks post-injection (Phillips, 1997). This results in a prolonged skin photosensitivity, an obvious disadvantage especially for patients with late stage malignancies. Nonetheless, the prolonged skin sensitivity induced by PDT with Photofrin[®] is only a mild inconvenience as compared to the severe adverse effects of standard chemotherapeutic regimens. While these disadvantages have not prevented Photofrin[®] from becoming a useful drug against cancer and other conditions, the search for new second generation photosensitizers with improved physical, chemical and spectral properties remains an important goal (Sharman et al., 1999; Ali and van Lier, 1999).

Ideal photosensitizers for PDT must meet certain criteria, (Bonnett, 1999; Bonnett, 1995; Wöhrle et al., 1998; van Lier, 1988; MacRoberts et al., 1989) namely:

- It should be chemically pure and maintain a constant composition throughout treatment, undergoing minimal photobleaching.
- It should have minimal dark toxicity.
- The photosensitizer should be preferentially retained by the target tissue, whether malignant tumour cells or viral components, so as to induce only marginal toxicity to surrounding healthy biological matter. In addition, the excess dye should be rapidly excreted from the body exhibiting low systemic toxicity.
- The dye should have high photochemical reactivity with a high quantum yield of long-lived triplet-states energetic enough to produce singlet oxygen.
- The sensitizer should have a large absorption coefficient at a long wavelength (600-800 nm) where there is optimal tissue penetration by light with a low degree of

attenuation by hemoglobin. Furthermore, cheaper diode lasers can be used in this range, thus increasing the potential utility of PDT in a clinical setting.

While no photosensitizer can be expected to fulfill all of these parameters, numerous second generation photosensitizers have been investigated that can overcome the shortcomings of hematoporphyrin derivatives and take advantage of their more highly suited properties in order to treat a number of conditions. For instance, the use of 5-aminolaevulinic acid (ALA)-induced endogenous photosensitizers has gained clinical acceptance for the treatment of actinic keratoses of the face and scalp. In the meanwhile verteporfin (benzoporphyrin derivative monoacid ring A) has received market approval in several countries including Switzerland, the United States and Canada, under the trade name Visudyne[®] as a therapy for wet age-related macular degeneration (AMD) a leading cause of blindness in people over the age of 50. Overall, the photosensitizer used to treat a given condition will greatly depend on the characteristics of the compound and the condition to be treated, with a wide range of different photosensitizers being used in clinic, each for its own particular use. Appendix II outlines the applications of several photosensitizers in pre-clinical and clinical trials.

1.4 Second generation photosensitizers: Phthalocyanines

Among the more promising second generation photosensitizers are phthalocyanines. The term phthalocyanine finds its origin in the Greek word “naptha” which means rock oil and “cyanine” which means dark blue. It was first used to describe this class of macrocyclic compounds by Sir Reginald Linstead in 1933 during his pioneering work on the subject (Linstead, 1934). In addition to the immediate applications as dyes and colorants, it was realized that such compounds would be of great

academic interest. As such, starting in 1929, Linstead and his group began their work on phthalocyanines which lead to the determination of their structure in the early 1930's (Linstead and Lowe, 1934; Dent et al., 1934; Linstead and Robertson, 1936). The synthesis of Pcs and their relationship to porphyrins, along with further examination into the intricate structure of Pcs, their planar nature, their complexes with metal ions and their stability were subsequently studied (Byrne et al., 1934; Linstead and Lowe, 1934; Dent and Linstead, 1934; Dent, 1938; Barrett et al., 1936; Barrett et al., 1938). The structure of phthalocyanines was later confirmed by Roberston via x-ray crystallography in a series of classic papers (Robertson, 1935; Robertson, 1936; Robertson and Woodard, 1937; Roberston and Woodard, 1940). In the seventy years to follow this initial work, a plethora of information emerged concerning diverse synthetic routes, Pc photochemical and photophysical properties and potential applications.

Of the utmost importance is their potential role in photodynamic therapy. These azaporphyrin derivatives have stronger absorbances at longer wavelengths than do porphyrins. Pcs have favorable photophysical and chemical properties. In addition, these properties can be altered through the addition of substituents to the periphery of the macrocycle or axial ligands to the chelated central metal ion. This makes them interesting photosensitizing agents.

Phthalocyanine derivatives have attractive photophysical and photochemical properties as compared to hematoporphyrin derivatives. Monomeric Pcs have a characteristic absorption spectra with a strong Soret absorption peak at approximately 350 nm, a weak maximum around 600 nm and a narrow, very strong, absorption peak in the far red region of the visible spectra ($\text{Pc } \lambda_{\text{max}} \approx 680 \text{ nm}$; $\text{HpD } \lambda_{\text{max}} \approx 630 \text{ nm}$) where

tissue penetration by visible light is superior. In addition to the red-shift of phthalocyanine Q band absorption maxima, Pcs have an improved capacity to absorb light, by two orders of magnitude over that of the highest Q band absorption of HpDs (Pc molar extinction coefficient (ϵ) in the $10^5 \text{ M}^{-1}\text{cm}^{-1}$ range; HpD $\epsilon \sim 10^3 \text{ M}^{-1}\text{cm}^{-1}$) (van Lier et al., 1988; van Lier and Spikes, 1989). These qualities therefore lead to enhanced photophysical and photochemical properties.

The nature of the central metal ion influences the Pc photophysical properties (triplet quantum yield and lifetime). Complexation of Pcs with open shell or paramagnetic metal ions such as Cu^{2+} , Co^{2+} , Fe^{2+} , Ni^{2+} , VO^{2+} , Cr^{3+} and Pd^{2+} give dyes with shortened triplet lifetimes (nanosecond range) due to increased intersystem crossing back to the ground state, which renders the dye photoinactive (Chan et al., 1987). Pcs containing closed d shell or diamagnetic metal ions, such as Zn^{2+} , Al^{3+} and Ga^{3+} , are dyes producing high triplet yields ($\phi_T > 0.4$) with long lifetimes ($\tau_T > 200 \mu\text{s}$) (Darwent, 1982). These triplet M-Pcs vary in energy from 110-126 kJ/mol which is ample energy to generate singlet oxygen (94.5 kJ/mol) with high quantum yields (ϕ) of approximately 0.3-0.5 (Wagner et al., 1987). Singlet oxygen yields are affected by ring substituents and axial ligands. Phthalocyanines tend to aggregate or dimerize which in turn determines the MPcS_n activity (Wagner et al., 1987). Aggregation is easily detected spectroscopically as absorption occurs at shorter wavelengths in the Soret region and there is a 30-50 nm blue-shift of the Q band absorption peak which appears more broad and less intense. Disaggregation to form monomeric dyes can be accomplished by adding detergents, serum or plasma proteins or organic solvents (van Lier and Spikes, 1989; Ben Hur et al., 1987).

1.4.1 Phthalocyanine induced apoptosis or necrosis

Phthalocyanines are very potent photosensitizers for use in PDT. Since first reported, several differently substituted Pc molecules have been synthesized and their PDT potential evaluated. Preliminary PDT results were promising using Pc derivatives, thus prompting more in depth studies regarding mode of cell death.

Apoptosis is a notable mode of cell death in response to PDT using phthalocyanines (Oleinick, 1998). A promising phthalocyanine receiving a great deal of attention is the silicon Pc, denoted as Pc4, bearing a long chain amino axial ligand ($\text{HOSiPcOSi}(\text{CH}_3)_2(\text{CH}_2)_3\text{N}(\text{CH}_3)_2$). This silicon phthalocyanine has shown promising results both *in vitro* and *in vivo*. Axial ligated Pcs have an important advantage over Pcs substituted on the periphery. Not only does the axial ligand impart greater solubility and prevent aggregation, these compounds do not have isomers, thus simplifying the preparation of pure samples.

Pc4 has been shown to initiate an apoptotic response in several cell lines including L5178Y-R cells. DNA fragmentation, a hallmark of apoptosis, was observed (He et al., 1997). This same research team demonstrated the importance of cell membrane damage with respect to apoptosis induction. Photodamaging the cell membrane initiates a cascade of events such as phospholipase C activation, release of secondary messengers and an increase in intracellular calcium concentration, all leading to cell suicide (Agarwal et al., 1993). Furthermore, an increase in intracellular calcium concentration also leads to the formation of nitric oxide (NO), an important bioactive signaling molecule. It has been demonstrated that NO is involved in PDT-mediated apoptosis (Gupta et al., 1998). In addition to NO, other well documented apoptotic

pathways are initiated by Pc4-PDT. Cytochrome c is released into the cytosol and subsequent activation of proteases, (Varnes et al., 1999) now known as caspases, resulting in the cleavage of various proteins including poly(ADP-ribose)polymerase have been documented (He et al., 1998). Cell cycle arrest in the G0/G1 phase of the cell cycle as a result of p21 induction has been observed following Pc4 therapy (Agarwal et al., 1993). In addition, the participation of retinoblastoma protein has been studied following phototherapy (Ahmad et al., 1998). Several other apoptotic pathways and proteins involved in PDT mediated apoptosis are under investigation.

Furthermore, enhanced apoptotic responses to photodynamic therapy with ALPc was observed after bcl-2 transfection (Choi Kim et al., 1999). Bcl-2 is a known inhibitor of apoptosis via antagonizing the release of mitochondrial cytochrome c. In this case, using immortalized MCF10A cells, overexpression of bcl-2 and bax was revealed and PDT caused selective destruction of bcl-2, leaving bax unaffected. As such, the greater apoptotic response towards PDT in this case is probably attributed to the higher bax:bcl-2 ratio after photodamage. On the other hand, CHO-K1 cells transfected with Bcl-2 exhibited a partial resistance to PDT-induced cell death, either via apoptosis or non-apoptotic mechanisms (He et al., 1996).

Subsequently, it was demonstrated that the mode of cell death is not only dependant on cell type (i.e. fibroblasts do not die via organized DNA fragmentation (He et al., 1997)) but also on the photosensitizer, its intracellular localization and the drug and energy dose administered. In addition, it has recently been shown that the mode of cell death also varies with cell density due to the bystander effect (Dahle et al., 1999). Using ALPc and lower PDT doses resulted in the induction of apoptosis as opposed to necrosis.

Organelles such as the mitochondria and lysosomes were mainly affected. At higher PDT doses, cells underwent severe membrane damage resulting in necrosis (Luo and Kessel, 1997). Interestingly, Dellinger reported a mixed response when using Photofrin® and CV-1 cells (Dellinger, 1996). At higher irradiation doses, the cells lysed instead of exhibiting fragmentation associated with apoptosis. However, morphological changes typical of apoptosis were observed initially, such as plasma membrane blebbing, condensation of the cytoplasm and nuclear fragmentation. These cells appear to enter an apoptosis-like death, yet this pathway is stopped by plasma membrane lysis and ultimately, the cells undergo necrosis. *In vivo* studies using the murine MS-2 fibrosarcoma tumor model demonstrated random necrosis of tumor cells post ZnPc-PDT. Interestingly, apoptotic pathways were also detected, suggesting different localization of photosensitizers within the cell upon illumination (Jori and Fabris, 1998).

Photodynamic therapy using phthalocyanines is known to induce a number of responses that ultimately lead to apoptosis. For instance, PDT using Pc4 induces an oxidative stress associated with increased ceramide generation (Separovic et al., 1997), which has been associated with apoptosis in several malignant and non-malignant cell lines. The increased production of ceramide is due to the stress-induced activation of sphingomyelinase (SMase). It has been shown that normal human lymphoblasts accumulate ceramide and undergo apoptosis following Pc4-PDT. In contrast, Niemann-Pick disease lymphoblasts, which are deficient in acid sphingomyelinase, fail to respond to Pc4-PDT via ceramide accumulation and apoptosis (Separovic et al., 1999). However, addition of exogenous bacterial Smase during Pc4-PDT induced a significant apoptotic response. This led to the hypothesis that SMase may be an important proapoptotic

factor determining responsiveness of cells to Pc4-PDT. Previous work showed that Pc4-PDT was able to induce apoptosis in numerous cell lines. Three cell lines were studied *in vitro* revealing the accumulation of ceramide post PDT and induction of apoptosis. Interestingly, there were increased levels of ceramide in RIF-1 cells post PDT with no evidence of apoptosis. Ceramide clearly is associated with PDT-mediated cell death either by inducing apoptosis or necrosis (Separovic et al., 1998).

1.4.2 Phthalocyanine delivery systems

Targeting of photosensitizers is imperative, as singlet oxygen is extremely reactive thus limiting PDT action to the site of singlet oxygen generation. As previously stated, its lifetime in biological systems has been estimated to be 200 ns with a diffusion range of 45 nm (Ochsner, 1997). Mammalian cells are 10^4 to 10^5 nm in diameter thus the importance of PS distribution within the target tissue is evident. Hydrophobic sensitizers have been shown to be biologically beneficial as compared to hydrophilic compounds. The former exhibits greater photocytotoxicity and tumor eradication, most probably because these hydrophobic dyes localize in more photosensitive cellular compartments. However in aqueous solutions, photosensitizers, either non-substituted or substituted with hydrophilic groups, tend to aggregate and dimerize which affects their cell penetrating properties and results in a loss of photochemical activity. Increased photodynamic action has been attributed to increased monomerization of the photosensitizer.

Lipophilic photosensitizers require biologically compatible delivery systems. Nanoparticles have received attention as potential delivery vehicles for drug targeting. A hexadecafluoro zinc phthalocyanine was formulated in polyethylene-glycol-coated poly(lactic acid) nanoparticles and evaluated for PDT efficiency. Using the EMT-6

tumor model, improved tumor response was observed using the nanoparticle formulation as compared to the same dye solubilized in Cremophor EL (Allémann et al., 1996). Nanoparticle formulation allows for increased serum levels of the dye as compared to Cremophor EL (CRM) formulation. Increased blood levels and enhanced tumor uptake lead to higher tumor-to-organ concentration ratios enhancing the overall dye efficacy (Allémann et al., 1995). Likewise pH-responsive, polymeric micelles consisting of random co-polymers of *N*-isopropylacrylamide, methacrylic acid and octadecyl acrylate have been loaded with water-insoluble AlPc and shown to exhibit higher phototoxicity against EMT-6 cells *in vitro* than the control CRM preparation (Taillerfer et al., 2000).

In addition to emulsifiers such as Cremophor EL, Solutol HS or lipid-based vehicles such as liposomes, a substantial amount of research has focused on proteinic carriers for improved targeting of malignant cells. Serum lipoproteins can incorporate hydrophobic sensitizers into their lipid portion. Frequently, neoplastic cells have increased LDL receptors on the cell surface as increased proliferation requires cholesterol for membrane synthesis (Vitols, 1991). Subsequently, LDL are known to deliver photosensitizers to neoplastic cells through a receptor internalization mechanism (Hamblin and Newman, 1994; Mazière et al., 1991; Allison et al., 1994). Jori and associates have done extensive studies using ZnPc incorporated into liposomes of varying composition. As will be discussed in greater detail later, Jori et al. have investigated the role of serum lipoproteins and found them to be effective delivery vehicles for phthalocyanines (Polo et al., 1995; Reddi, 1997; Reddi et al., 1990; Valduga et al., 1999).

Recently, a series of publications has reported the use of a modified aluminum tetrasulfophthalocyanine covalently coupled to various proteins via one or two caproic

acid spacer chains. Brasseur et. al. targeted the scavenger receptor of macrophages using a maleylated bovine serum albumin covalently labeled with AlPcS₄. Relative photocytotoxicities were reported using a receptor positive and a receptor negative cell line where their lethal effects correlated with receptor affinity for the BSA-Pc conjugate (Brasseur et al., 1999).

Using the same phthalocyanine derivative, it was found that covalently labeling a monoclonal antibody for the CD3+ antigen in a similar manner did not hinder receptor recognition. There was increased uptake of the antibody-Pc conjugate in a receptor positive cell line as compared to the free phthalocyanine and antibody-Pc uptake was less in a cell line void of receptors (Ménard, 1998). However, the Pc-antibody conjugates were photoinactive, most likely due to inefficient subcellular targeting. Similarly, AlPcS₄ was covalently coupled to monoclonal antibodies directed against carcinoembryonic antigen (CEA) (Carcenac et al., 1999). *In vivo* studies using nude mice bearing human colon carcinoma xenografts (T380) demonstrated increased uptake of the antibody-Pc conjugate as compared to the free Pc. *In vitro* results were encouraging using this antibody complex.

1.5 Research objectives

Over the past 20 years, several photosensitizers have been synthesized and tested both *in vitro* and *in vivo* with varying results. Initially, this approach to PDT was sufficient. However, more elaborate research is now underway to improve efficiency by targeting the photosensitizer (PS) to various tissue types. A number of second generation photosensitizers such as phthalocyanines are effective drugs for PDT. However, to improve tumor uptake and preferential tumor-to-normal tissue ratios, third generation photosensitizers are being designed where the chromophore is tagged with a targeting system in order to improve these characteristics.

The current research is sub-divided into four categories as represented by four research papers that have been published or submitted for publication as part of the present study. Chapter 2 presents results of a study employing a series of AlPc derivatives substituted with aliphatic chains of varying lengths on the Pc macrocycle imparting different lipophilicity. Amphiphilicity plays a major role in determining the usefulness of a Pc for PDT. The first study investigates the PDT potential of a series of amphiphilic AlPc derivatives as a function of lipophilicity. This study tests the hypothesis that amphiphilic phthalocyanines have improved PDT potential.

Subsequently, the role of low-density lipoproteins (LDL) as vehicles for AlPcS delivery to target tumor cells was evaluated and reported in chapter 3. Parallel studies using both non-covalently labeled and covalently labeled LDL to target human adenocarcinoma lung cancer cells (A549) were carried out. The amphiphilic AlPcS₄ molecule bearing a twelve-carbon alkyl side chain on the Pc macrocycle was noncovalently inserted in the LDL phospholipid layers. AlPcS₄ was covalently coupled to

the apolipoprotein moiety via two sulfonamide five-carbon spacer chains. The two fold question addressed was whether or not LDL molecules are suitable vehicles for Pc delivery to the target tissue and whether Pc attachment to the protein or the lipid portion of the LDL affects PDT.

Using the same principle, adenoviral proteins were employed to target lung cancer cells rich in the appropriate class of integrin receptors. Adenoviruses have been receiving a great deal of attention as gene therapy vectors. These viral particles gain entrance into the cells via receptor mediated endocytosis. Two receptors are required, the first for attachment and the other mediating internalization. Internalization of the virus particle is via integrin receptors able to bind with high affinity to the RGD binding motif found in the penton base protein of adenovirus. Chapter 4 outlines studies using the adenovirus capsid proteins covalently labeled with ALPcS₄ derivatives. Is Pc delivery to cells improved and is PDT more efficient as a result of this?

Tumor targeting using protein vehicles may have serious limitations invoking adverse immune responses. The search for small peptidic vectors was the rational behind the fourth objective of this thesis as described in chapter 5. The RGD peptide sequence is currently being investigated as a targeting vehicle for conventional chemotherapeutics. The present study evaluated the effects of RGD on cell survival following ALPcS_{2adj} photodynamic treatment of both a receptor negative and a receptor positive cell line. Does RGD enhance Pc-PDT?

Chapter 2.

Photodynamic properties of amphiphilic derivatives of aluminum tetrasulfophthalocyanine

Cynthia M. Allen, Réjean Langlois, Wesley M. Sharman, Carole La Madeleine and Johan E. van Lier (2002) *Photochem. Photobiol.* **76**, 208-216.

Pour l'article complet voir la copie papier à la Bibliothèque des sciences de la santé
Section Monographie WB 480 A57 2001

Chapter 3.

Low Density Lipoprotein-Bound Aluminum Sulfophthalocyanine: Targeting Tumor Cells for Photodynamic Therapy

Pascale Urizzi, Cynthia M. Allen, Réjean Langlois, René Ouellet, Carole La Madeleine and Johan E. van Lier (2001) *J. Porphyrins and Phthalocyanines* **5**, 154-160.

Pour l'article complet voir la copie papier à la Bibliothèque des sciences de la santé
Section Monographie WB 480 A57 2001

Chapter 4.

Symposium-in-Print

Photodynamic therapy: Tumor targeting with adenoviral proteins

Cynthia M. Allen, Wesley M. Sharman, Carole La Madeleine, Joseph M. Weber, Réjean Langlois, René Ouellet and Johan E. van Lier (1999) *Photochem. Photobiol.* **70**, 512-523.

Pour l'article complet voir la copie papier à la Bibliothèque des sciences de la santé
Section Monographie WB 480 A57 2001

Chapter 5.

Attenuation of photodynamically induced apoptosis by an RGD containing peptide

Cynthia M. Allen, Wesley M. Sharman, Carole La Madeleine, Johan E. van Lier and Joseph M. Weber (2002) *Photochem. Photobiol. Sci.* **1**, 246-254.

Pour l'article complet voir la copie papier à la Bibliothèque des sciences de la santé
Section Monographie WB 480 A57 2001

Chapter 6.

Discussion

6. Discussion

Second generation photosensitizers have several advantages over hematoporphyrin derivatives. However even the most potent chemotherapeutic agents are of no use if they don't target the desired tissue and can have deleterious effects. This alone outlines the importance of drug targeting.

6.1 Amphiphilic phthalocyanines

Substituents on the phthalocyanine framework have included aliphatic chains and higher order aromatics to improve both the solubility and potential usefulness of the Pc. The current study uses AlPcS₄ substituted with aliphatic chains of increasing length. In general, amphiphilic molecules have been found to be more photodynamically active. The hydrophobic chain portion can slip into the cell membrane and anchor the Pc while their charged sulfonate groups interact with the aqueous layer outside the cell. Following 48 hours incubation, the highest cellular concentration was observed with the more lipophilic AlPcS₄(C16) derivative followed by the C12 < C8 < C4. The more hydrophilic compounds accumulated to a less extent. The AlPcS_{2adj}, having a similar retention time on the HPLC column as the AlPcS₄(C12), also had comparable A549 cell uptake properties. It is important to stipulate cell type as different cell types may exhibit different membrane permeation of the different Pc molecules. In addition, decreasing aggregation of the phthalocyanine enhances cell uptake of the sensitizer. In the present study the AlPcS₄(C16) was initially solubilized in methanol then further diluted in PBS. The methanol was evaporated off leaving only trace amounts. Studies where the AlPcS₄(C16) derivative was dissolved directly in PBS resulted in different uptake by the

cells of the photosensitizer and the phototoxicity of the sensitizer was altered. This is most likely due to increased sensitizer aggregation.

To circumvent such problems as aggregation but to retain the amphiphilic nature of the sensitizer, a series of silicon phthalocyanines with charge axial ligands was synthesized and tested on the EMT-6 cell line (Murphy et al., 2001). Axial ligated Pcs have an important advantage over Pcs substituted on the periphery. Not only does the axial ligand impart greater solubility and prevent aggregation these compounds do not have isomers, thus simplifying the preparation of pure samples. The SiPc in this study, with the longest axial ligand thus the more amphiphilic, was most phototoxic. This is comparable to the present study where the $\text{AlPcS}_4(\text{C16})$ molecule was most photoactive. This increased length of the axial substituent presumably helps the dye to better partition into the membrane. A promising amphiphilic phthalocyanine receiving a great deal of attention is the silicon Pc, denoted as Pc4, bearing a long chain amino axial ligand $(\text{HOSiPcOSi}(\text{CH}_3)_2(\text{CH}_2)_3\text{N}(\text{CH}_3)_2)$. Clinical trials are scheduled to begin in 2001 (personal communication).

6.2 Protein carriers

Site-specific drug delivery is an integrated part of the intellectual and experimental design of a therapeutic system. Enhanced delivery systems can access previously inaccessible cellular compartments thus reducing the amount of drug administered and protecting the body from unwanted accumulation and toxicity. The dye must be exclusively delivered to the pharmacological site of action at an appropriate dose to effectively combat the disease and improve the therapeutic index. As the pathophysiology of various diseases becomes known, researchers can exploit cellular

systems to increase dye uptake. Receptor-mediated endocytosis allows for the uptake and intracellular processing of macromolecules that bind to the cell surface receptors. Endocytosis of the plasma membrane is a continuous activity in many cell types occurring at a constant rate regardless of external stimuli. In this way, cell surface receptors are regularly re-distributed to the plasma membrane. Receptor internalization can also be stimulated by ligand binding. It has been estimated that 80% of the epidermal growth factor (EGF) receptors internalize within a half time of 2.5 minutes upon binding saturation (Hopkins, 1986). In view of this, drug trafficking across the cell plasma membrane is feasible. Additionally, protein carriers specifically targeting a cellular population will decrease liver entrapment of the photosensitizers as the drug package is strictly addressed. Several protein/receptor systems have been explored for the purpose of drug delivery.

We report that the presence of 4 $\mu\text{g/mL}$ of LDL slightly enhances the phototoxicity of the AlPcS₄ derivatives substituted with long chains. Naturally, the next phase in this study was to incorporate these lipophilic sensitizers with LDL prior to administration. This was accomplished either via covalent or non-covalent coupling of the Pc to the LDL. Extensive studies have investigated serum proteins as drug carriers and merit acknowledging.

6.3 Serum proteins

Serum proteins are predominantly responsible for the transport of photosensitizers throughout the body. Serum albumin generally transports hydrophilic PS whereas more hydrophobic PS localize preferentially in lipoproteins, leading to enhanced intracellular accumulation of the dye via receptor mediated endocytosis (Table 6.1) (Rosenthal, 1991;

Reddi, 1997; Hamblin and Newman, 1994a). To further benefit from this, coupling of photosensitizers with albumin and low-density lipoproteins (LDL) prior to administration has been investigated as a means to deliver PS with higher target specificity.

6.3.1 Albumin

Initial studies involved the non-covalent binding of PS to bovine serum albumin (BSA) prior to intravenous administration (Larrocque et al., 1996). Using two *in vivo* tumor models, murine mammary carcinoma and human colon carcinoma, tumor regression was observed at ZnPc doses of 0.5 and 2.0 μmol per kg respectively.

Alternatively, covalent conjugates were prepared using BSA and various photosensitizers. These biomolecules can enter the cell either via passive diffusion or via receptor mediated endocytosis. Tumor-associated macrophages accumulate higher levels of photosensitizer as compared to neighboring tumor cells (a nine fold increase in the case of porphyrins) (Hamblin and Newman, 1994b). It has been estimated that in several cancers over 50% of the tumor mass is of the macrophage lineage. Macrophages express high numbers of the scavenger receptor which is able to bind a wide range of ligands, shuttling them to endosomes and lysosomal compartments. This receptor readily binds oxidized LDL and maleylated bovine serum albumin but not the corresponding native proteins (Krieger et al., 1993). Thus by covalently attaching photosensitizers to the BSA molecule, specific targeting of this receptor ensues.

Brasseur et al. reported that maleylated-BSA was covalently coupled to AlPcS_4 via one or two sulfonamide hexanoic amide spacer chains. The photosensitizers used in the preparation of these bioconjugates is the same as that used in the covalent labeling of the adenoviral proteins (chapter 4) and the LDL molecules (chapter 3). The Pc-conjugate

was readily taken up by the scavenger receptors on J774 cells. Mal-BSA-Pc was more photocytotoxic against this receptor-bearing cell line compared to a negative control (Brasseur et al., 1999). Similarly, mal-BSA coupled to chlorin e6 exhibited increased PS accumulation and phototoxicity when using J774 cells as compared to non-target OVCAR-5 cells (Hamblin et al., 2000a; 2000b). Dye uptake was decreased after incubating the mal-BSA-Ce6 with the cells at 4°C, indicating an endocytotic route of entry.

6.3.2 Lipoproteins

The current study evaluates low density lipoproteins (LDL) as potential dye carriers. LDL are particles that are naturally produced by the liver. Their primary role is to transport cholesterol and other lipids to various cells throughout the body. They possess a receptor-binding motif, apolipoprotein B-100, which binds with high affinity to a LDL cell surface receptor and is subsequently internalized. Cancer cells naturally express greater numbers of these receptors, making LDL particles attractive vehicles for drug targeting (Gueddari et al., 1993).

Early *in vivo* studies showed that very hydrophobic photosensitizers could only be administered following inclusion in lipid material (Reddi et al., 1990). Therefore ZnPc was enclosed in various liposomes. Several studies have investigated the interactions of LDL and liposomes containing ZnPc to determine whether the ZnPc becomes incorporated in the LDL (Valduga et al., 1999; Versluis et al., 1994; Rensen et al., 1994). Following *in vitro* incubation of the LDL and liposomes, the LDL particles acquired a net negative charge as seen using electrophoretic mobilities. This is indicative of ZnPc loading into the LDL. Consequently, intravenous injection of liposomal Pc showed that

LDL particles were indeed involved in Pc transport. Therefore, studies in which the ZnPc was non-covalently complexed with LDL *prior* to injection showed a two-fold greater tumor uptake of the Pc than when the Pc was enclosed in DPPC liposomes (Reddi et al., 1990). Accordingly, the use of the LDL receptor pathway as a tool for enhancing the selectivity of PS delivery to tumor cells has been extensively studied (Mazière et al., 1991; Jori et al., 1993).

AlPc was non-covalently incorporated into oxidized LDL and the photodynamic efficiency evaluated. The oxidized LDL-Pc complex was taken up by the scavenger receptor after incubation with macrophages. Incubation of this Pc-LDL complex in serum revealed that it was quite stable, showing very little re-distribution amongst the other serum components (de Vries et al., 1999). In addition, LDL were non-covalently labeled with hematoporphyrin. Selectivity of tumor targeting was prominent. Electron microscopy demonstrated that the Hp-LDL conjugate induced direct tumor cell kill as opposed to the indirect effect seen when free Hp is administered. This suggested that targeting was through the LDL receptor pathway as the Hp-LDL seemed to be taken up by the cells via receptor-mediated endocytosis (Zhou et al., 1988).

Chapter 3 outlines the study whereby AlPcS₄ with a twelve carbon alkyl chain (C₁₂) was non-covalently inserted into the phospholipid moiety of the LDL (Figure 3.1). *In vitro* PDT studies on adenocarcinoma lung cancer cells (A549 cells), which have increased LDL receptor expression, using the AlPcS₄C₁₂ conjugate were very effective (Urizzi et al., 2001). The AlPcS₄C₁₂-LDL conjugate was twice as phototoxic as compared to the AlPcS₄C₁₂, whereas, under similar experimental conditions, the parent molecule, AlPcS₄, was inactive. The AlPcS₄C₁₂-LDL bioconjugate was more

photoactive against the EMT-6 cell line. At doses as low as $0.2 \mu\text{mol kg}^{-1}$, tumor regression was observed using the bioconjugate against the murine EMT-6 model. However as shown in chapter 2, the AlPcS₄ derivatives substituted with long side chains exhibit similar tumor response and did not require coupling of the Pc with LDL prior to administration.

While non-covalent labeling of LDL particles showed promise, attempts to make covalent conjugates have been disappointing. LDL were covalently coupled with AlPcS₄ bearing two sulfonamide five carbon chains (A₂) (Figure 3.1). Covalent coupling of the LDL protein moiety via the carboxylic acid groups of the monocaproic acid substituents reduced receptor recognition, thus decreasing cell/LDL-Pc interactions. Molar Pc:LDL ratios of approximately 70:1 were maintained as this represented 90% labeling yield. Perhaps decreasing this ratio would enhance cell uptake of the bioconjugate leading to increased phototoxicity. This Pc conjugate was photo-inactive at the highest drug and light doses studied. An alternative explanation could be altered trafficking upon internalization, without redistribution of the Pc to more photo-susceptible sites within the cell. Likewise, Hamblin and Newman conjugated LDL with hematoporphyrin via an amide bond which is similar to the present study (Hamblin and Newman, 1994c). Confocal microscopy showed that Hp-LDL localized in lysosomal compartments and that the LDL receptor was not solely responsible for uptake of this conjugate. Phagocytosis played a key role as well.

Alternatively, other photosensitizers have been covalently attached to LDL via carbodiimide activation such as chlorin e6 (Schmidt-Erfurth et al., 1997). LDL receptors on retinoblastoma cells were targeted. There was a 5-fold increase in chlorin e6 uptake

when using the LDL-Ce6 conjugate as compared to free Ce6. Similarly, the LDL-Ce6 complex was very phototoxic. These results suggest that LDL could be used as dye carriers for intraocular tumors such as retinoblastomas thus avoiding invasive treatments.

BPD non-covalently associated with LDL was likewise used to treat ocular conditions. It was observed that BPD-LDL accumulation was insignificant in LDL receptor negative fibroblasts or when the LDL was chemically modified. Furthermore, *in vivo* studies in M1 tumor bearing DBA/2J mice showed a pronounced enhancement of BPD-native LDL tumor accumulation over that of acetylated LDL associated BPD (Allison et al., 1994). These results indicate that the LDL receptor is responsible for the uptake of the BPD-LDL. Subsequently, BPD-LDL photodynamic therapy was tested on an experimental model of choroidal melanoma (Schmidt-Erfurth et al., 1994), experimental choroidal neovasculation (Miller et al., 1995) and Greene melanoma (Schmidt-Erfurth et al., 1996). Results were favorable in all three models indicating LDL usefulness as a sensitizer delivery vehicle.

Table 6.1 Summary of photosensitizer serum-based vehicles and bioconjugates

Vehicle	Photosensitizer	Target	Reference
Albumin (BSA)	Hematoporphyrin	Macrophages	Hamblin and Newman 1994
Albumin (BSA)	ZnPc	Murine mammary and human colon carcinoma	Larroque et al. 1996
Albumin (BSA), Fibrinogen, Gelatin	Chlorin e6	Tissue solder	Khadem et al., 1999
Maleylated BSA	Chlorin e6	Scavenger receptor	Nagae et al. 1998
Maleylated BSA	AlPcS ₄ A ₁ and A ₂	Intimal hyperplasia Scavenger receptor	Brasseur et al. 1999
BSA and Maleylated BSA	Chlorin e6	Scavenger receptor	Hamblin et al. 2000
LDL	ZnPc	MS-2 fibrosarcoma	Reddi et al 1990
LDL	TPPS _n	LDL receptor on human hepatocyte tumor	de Smidt et al 1993
LDL (human)	BPD-MA	Choroidal melanomas, choroidal neovasculture	Schmidt-Erfurth et al 1994 Miller et al., 1995
LDL, HDL	Hematoporphyrin	LDL receptor on fibroblast Scavenger receptor	Hamblin and Newman, 1994 Hamblin and Newman, 1994
LDL	BPD-MA	LDL receptor on rhabdomyosarcoma	Allison et al 1994
LDL (human)	BPD-MA	Greene melanoma	Schmidt-Erfurth et al 1996
LDL (human)	Chlorin e6	Retinoblastoma	Schmidt-Erfurth et al 1997
LDL (oxidized)	AlPc	Scavenger receptor	Devries et al 1999
LDL	AlPcS ₄ A ₂ and AlPcS ₄ (C ₁₂)	LDL receptor of lung adenocarcinoma	Urizzi et al 2000
Transferrin	Hematoporphyrin	Transferrin receptor	Hamblin et al. 1994

6.4 Antibody conjugates

A class of protein vehicles is the antibody vehicles. Photodynamic therapy using PS-Ab conjugates, termed photoimmunotherapy (PIT), has been developed over the past 15 years (Hasan et al., 1997; Klyashchitsky et al., 1994). PIT has evolved in parallel with the use of monoclonal antibodies as pharmaceuticals. PIT is an ever-expanding field of study as mAb generation techniques improve thus generating different antibodies to be coupled with various photosensitizers. Targeting of photosensitizers using antibodies and/or fragments has been extensively investigated. Table 6.2 is a summary of various studies.

6.4.1 Phthalocyanines

Phthalocyanines have likewise been targeted to various cell lines via antibody carriers. AlPcS_n was encapsulated in liposomes that were then linked to antibodies for targeting purposes. It was clear from these early studies that the phototoxicity was directed by the mAb because free Pc or liposomal Pc were not phototoxic against the antigen-bearing cell lines under the same conditions (Morgan et al., 1989). Subsequently, this liposomal AlPcS_n formulation was used to target B-cells using an anti-B-cell Ab as well as T-lymphocytes using anti-CD3 antibody. This could be applicable for *ex vivo* bone marrow purging (Morgan et al., 1992). A similar method has also been used to target human bladder carcinoma (Morgan et al., 1994). An AlPcS_4A_1 (A_1 being a 5 carbon spacer chain) has been used to label anti-carcinoembryonic antigen (CEA) monoclonal antibodies. This is the same Pc derivatives used to label proteins in the current study. The Pc-mAb conjugates had 5-16 moles of Pc per mol of antibody. The immunoreactivity of the bioconjugates was not altered due to the coupling with a

photosensitizer. In nude mice, tumor to normal tissue ratios were similar for the Pc-mAb conjugates as compared to the free antibody. *In vitro*, the bioconjugate was photocytotoxic against antigen positive cells. This, taken together with the *in vivo* tumor seeking capacity of the Pc-mAb, makes it a promising third generation photosensitizer (Carcenac et al., 1999). Conversely, an anti-CD3 antibody was covalently labeled with AlPcS₄A₁. Cell uptake of the bioconjugate was not impaired as measured by flow cytometry. However, the bioconjugate was not photoactive (Ménard, 1998). It was postulated that upon internalization of the Pc-mAb, the photosensitizer was trapped in the endosome and following illumination, the singlet oxygen produced was quenched by the Pc-mAb microenvironment.

Table 6.2 Summary of photosensitizer-antibody conjugates

Antibody	Photosensitizer	Target	Reference
Rabbit anti-Hu CAMAL-1 L1210	Hematoporphyrin	Human peripheral blood leukocytes CAMAL antigen Irrelevant MoAb	Mew et al., 1985
Anti-Leu/1	Chlorin e6	T-leukemia cells	Oseroff et al., 1986
Mab 3G2/C6	Chlorin e6	Bladder carcinoma	Hasan et al., 1989
Mab 791T/36	Liposomal ALPcS	Osteosarcoma, colorectal carcinoma	Morgan et al., 1989
Mab 3G9	Hematoporphyrin	Gastric cancer	Xu et al., 1989
Mab 3H11	derivative		
Mab T48	BPD-MA	Human chorionogonadotropic (HCG) hormone	Jiang et al., 1990
Rabbit IgG anti- E.coli	Iodinated fluorescein	E. coli	Devanathan et al., 1990
Mab OC125	Chlorin e6	Ovarian carcinoma	Goff et al., 1991, 1994, 1996 Hamblin et al., 1996 Prosser et al., 1991
Mab HMFG-1	Hematoporphyrin	Breast adenocarcinoma	
Mab E7	Liposomal ALPcS	Urinary bladder carcinoma	Morgan et al., 1994
Mab OC125	Chlorin e6	Human ovarian cancer	Duska et al., 1997
Mab OKT3	ALPcS ₄ A ₁	Leukemic T lymphocytes (CD3+)	Menard, 1998
Mab U36	m-THPC	Head and neck squamous cell carcinoma	Vrouenraets et al., 1999
Mab 35A7	ALPcS ₄ A ₁	Carcinoembryonic antigen (CEA)	Carcanac et al., 1999
Mab U36	TrisMPyP-	Squamous cell carcinoma	Vrouenraets et al., 2000
Mab 425	ΦCO ₂ H		
Murine Mab 17.1A	Chlorin e6	Human colorectal cancer antigen (EpCMA)	Del Governatore et al., 2000(a, b) Hamblin et al., 2000

6.5 Epidermal growth factor

Epidermal growth factor (EGF) is a small protein that binds to a cell surface receptor. EGF is a potent mitogen found throughout the body and is classified as an angiogenesis stimulating factor. As seen with integrin receptor expression, the EGF receptor (EGF-R) is over expressed in a variety of tumor types, in particular ovarian cancers. 30% of ovarian cancers have increased EGF receptor numbers, making EGF a potential drug carrier (Moor et al., 2000). EGF has been coupled to various photosensitizers in the hopes of specifically targeting this receptor. The cytotoxic activities of an AlPc and a CoPc bound to EGF were determined against a human breast carcinoma cell line. The AlPc-EGF was seven times more photoactive as compared to the free AlPc. The CoPc-EGF conjugate was cytotoxic, requiring ascorbic acid for activation as opposed to light. *In vivo* studies using B16 melanoma were subsequently carried out (Lutsenko et al., 1999).

More elaborate conjugates have been prepared using EGF and Sn(IV)chlorin e6 bound together via carriers such as dextran, polyvinyl alcohol and human serum albumin (HSA) (Gijssens and de Witte, 1998; Gijssens et al., 2000). The EGF-HSA-SnCe6 conjugate had the best receptor affinity displaying high phototoxicity against a cell line over-expressing the EGF-R. The effect was receptor related as native EGF could compete for binding sites and decreased photoactivity. The HSA conjugate also produced more intracellular ROS upon irradiation as compared to the dextran conjugate. In a previous study, the EGF-Dex-SnCe6 conjugate bound specifically to the EGF receptor on a squamous cell carcinoma line whereas the EGF-PVA-SnCe6 conjugate

displayed limited binding. It was concluded that the photodynamic activity of these conjugates was dependent on the carrier used.

6.6 Adenoviral Carriers

Adenoviral proteins were studied as protein carriers for ALPcS₄ (chapter 4). Adenoviruses have natural tropism for a variety of tissue types. Adenoviruses enter cells via receptor mediated endocytosis involving two separate receptors. The first one, designated as CAR (Coxsacki and Adenovirus Receptor), is responsible for mediating attachment while the second receptor is responsible for internalization. This second receptor is a member of the integrin class of surface receptors. As previously stated, integrin receptor expression is frequently altered in cancer cells. Studies have shown that human lung cancer cell lines such as A549 (used in the current study) have increased expression of RGD binding integrins on the cell surface (Hirasawa et al., 1994). This would suggest that adeno-vectors could function as selective carriers to deliver photosensitizers specifically to lung cancer cells.

Adenovirus type 2 capsid proteins were covalently labeled using an ALPcS₄ derivative via one or two caproic acid spacer chains (Figure 4.2). The specific adenovirus capsid proteins, the hexon, the penton base and the fiber were purified and conjugated with the Pc. The purification procedures were long and tedious resulting in only minimal amounts of purified proteins. Therefore, a mixture of adenovirus type 2 soluble proteins was also covalently coupled with ALPcS₄A₂. The bioconjugates were found to be only slightly photoactive *in vitro* when using the receptor positive cell line, A549 cells. *In vivo* studies were more promising. The Ad2-soluble protein mixture labeled with the ALPcS₄A₂ derivative, at doses of 0.5 $\mu\text{mol kg}^{-1}$ and 400 J cm⁻², resulted

in complete tumor regression using the EMT-6 model. This was a 20 fold decrease in Pc dose as compared to the parental AIPcS₄. Within the adenoviral protein mixture, the appropriate combination of viral proteins may be present invoking endosomal degradation to release the Pc. Endosomal entrapment of the photosensitizer has been proposed as an explanation for the decreased photocytotoxicity induced by the Pc conjugated with the purified penton base. This would prevent targeting of the PS to susceptible intracellular sites.

Increased cell uptake of the bioconjugate over that of the free Pc was seen, as would be expected due to the presence of integrin surface receptors that can readily bind the RGD motif of the penton base. However, there was only marginal cell kill after irradiation *in vitro*. Several factors are required to release the contents of the endosome upon internalization, in particular the adenoviral protease (Cotton and Weber, 1995). Therefore one could foresee using an attenuated adenovirus covalently labeled with a phthalocyanine as a delivery vehicle. The virus could be modified in such a way as to express one or more integrin binding ligands (RGD sequences) at the knob of the fiber protein. Upon binding via the RGD/integrin interaction, the virus-Pc would be internalized. The subsequent release from the endosome would be facilitated due to the use of an intact viral particle.

Sophisticated experiments were conducted using chlorin e6 as a photosensitizer, insulin as an internalizable ligand and the SV40 large T antigen as a nuclear localizing signal. These were all bound to BSA, a protein carrier. This conjugate was co-administered with an attenuated adenovirus (Akhlynina et al., 1999). Insulin/receptor complexes are internalized into endosomes and this is a limiting factor in the nuclear

transport of the conjugate as it limits the distribution of the sensitizer from locating in photosensitive sites. Adenoviruses aptly break open endosomes upon infection and therefore it was postulated that the bioconjugates would target the nucleus much more quickly and efficiently in the presence of adenoviruses. This indeed proved true. Rosenkranz and Sobolev have written comprehensive reviews on this subject (Rosenkranz et al., 2000; Sobolev et al., 2000).

Adenoviruses have yet another potential role in photodynamic therapy, that of a gene therapy vector. A yeast vector system for manipulating the adenoviral genome has been developed and used to construct a virus expressing a mutant form of ALA-synthase (Gagnebin et al., 1999). This mutated enzyme lacks the iron response element which regulates its translation and import to the mitochondria. This lead to the accumulation of endogenous photosensitizers in infected cells, conferring photosensitivity. Unlike conventional PDT this viral approach restricts photosensitivity by biological rather than purely physical or chemical means thus improving targeting.

6.7 Advances toward peptidic carriers

Small peptides are readily prepared by automated solid phase peptide synthesis. The several possible combinations of natural and non-natural amino acid sequences make this a versatile class of molecules. In addition peptidic vectors are desirable as they invoke minimal humoral and sytemic immune responses. The RGD motif binds with high affinity to several of the integrin class of cell surface receptors. As discussed in chapter 5, RGD peptides have several therapeutic applications. One such application was as a stimulator of apoptosis, in particular of endothelial cells of angiogenic vessels (Brooks et al., 1994). A second potential use is as a carrier for chemotherapeutic agents

such as doxorubicin (Arap et al., 1998). In the hopes of developing either a peptidic carrier system for phthalocyanines or an adjunct therapy to PDT, chapter 5 details a study of an RGD containing peptide used in conjunction with ALPc_{S2adj} PDT. It was found that a soluble RGD peptide attenuated PDT induced apoptosis. Generally, a soluble RGD peptide, when in contact with adherent cells disrupts the extracellular matrix thus blocking integrin signaling to the ECM and stimulates apoptosis. Buckley proposed an alternative method, that soluble RGD causes apoptosis via directly binding to and activating procaspase 3 (Buckley et al., 1999). At higher concentrations (10 mM) of the RGD peptide, we observed apoptosis with both a receptor positive and a receptor negative cell line therefore suggesting no involvement of the integrin receptor. At slightly inferior concentrations, there was a protection against PDT induce apoptosis in both cell lines with a much more pronounced protection in the receptor positive A549 cells. Not to completely rule out integrin receptor involvement, Whitlock and associates found that $\alpha_M\beta_2$ integrin signaling could either enhance or inhibit apoptosis of neutrophils depending on the proapoptotic stimuli (Whitlock et al., 2000). Activation of this class of integrins accelerated apoptosis induced by UV irradiation, Fas ligation and TNF α . However, integrin clustering without activation, resulted in inhibition of apoptosis.

In the current study, quenching of the reactive oxygen species by the peptide was ruled out as several different experimental parameters were changed to physically remove the peptide prior to PDT. The protective effect was still apparent and significant in the A549 cells whereas in the receptor negative cell line, EMT-6, the protection against apoptosis was almost abrogated. It was proposed that the RGD peptide, when rinsed, was

readily released from the EMT-6 cells, thus diminishing the protective effects whereas in the A549 cells, the peptide release was less. Studies using a radiolabeled peptide could prove useful in this capacity to determine peptide movement across the different cell membranes.

The A549 cells were previously shown in chapter 4 to bind with high affinity the radiolabeled RGD peptide whereas no significant binding was apparent in the EMT-6 cells. It is interesting that RGD has the most effect on A549 apoptosis. It could be possible that two different mechanisms are involved in the protective effect depending on the cell line. A549 cells, when incubated overnight with RGD prior to Pc-PDT, had enhanced cell survival whereas the EMT-6 had none. Perhaps, a receptor mediated, albeit a low affinity binding receptor, plays a role in the A549 protection whereas, simple diffusion of the peptide across the EMT-6 membrane may account for enhanced EMT-6 cell survival under certain circumstances. In the former, perhaps there is increased anti-apoptotic protein expression required to decrease cell death, whereas in the latter, perhaps the RGD is able to get into the cell and physically disrupt the apoptotic cascade.

Studies involving an immobilized RGD peptide could help elucidate whether cell penetration is required to abrogate apoptosis. In addition, it was postulated that increased expression of anti-oxidants such as superoxide dismutase, glutathione peroxidase and catalase may be a result of RGD stimulated integrin signaling (Qin et al., 2001). Clustering and activation of integrin receptors promotes *de novo* protein synthesis which was an absolute requirement to promote the survival effect of the extracellular matrix (Bozzo et al., 1997). Hence, growing cells on RGD coated plates, prior to PDT, may help to determine whether integrin signaling is involved in cell survival under these

circumstances. Analysis of protein levels could prove invaluable in elucidating the signals involved in RGD protection from PDT induce apoptosis.

6.8 Conclusion

Since its fortuitous beginning a century ago, photodynamic therapy has grown exponentially into a useful therapy for a number of conditions. The clinical acceptance of Photofrin®, methylene blue, Visudyne™ and the prodrug ALA clearly shows the extreme promise of PDT as a treatment for both malignant and non-malignant conditions for which there are few other useful therapies. There are a number of second generation photosensitizers in the pipeline with superior photophysical and biological properties. The choice of vehicles and carriers has dramatic effects on the properties of these photosensitizers. Clearly the investigation and development of new vehicles and 3rd generation photosensitizers could allow PDT to reach its full potential. There remains a long list of potential vehicles and carriers for photosensitizers including growth factors, hormones, steroid analogs, signaling molecules, peptides, antibody fragments, viral components, oligonucleotides and other tumor and tissue specific compounds.

The current study confirms that phthalocyanines are potent photosensitizers that are readily substituted, yielding better uptake and photodynamically active agents. Monosubstitution of AlPcS₄ with carbon chains of varying lengths increases the efficacy of this photosensitizer both *in vitro* and *in vivo*. L-Tryptophan oxidation following irradiation surpassed that of the parental AlPcS₄ molecule when the lipophilic sensitizers were monomerized in 1 % CRM. Cell adsorption as well as photocytotoxicity using the A549 cell line varied directly with the degree of hydrophobicity: C16 > C12 > C8 > C4. A549 cells took up 7 times more AlPcS₄(C16) as compared to AlPcS₄(C4) after a 48 hour

incubation. Human LDL at 4 $\mu\text{g/mL}$ decreased cell uptake of the Pc derivatives but had little or no effect on the cytotoxicity of the sensitizer presumably due to better subcellular localization. These lipophilic phthalocyanines were potent sensitizers *in vivo* using the EMT-6 tumor model. Complete tumor regression was obtained at doses as low as 0.2 $\mu\text{mol kg}^{-1}$ using both the C12 and the C16 derivatives whereas AlPcS₄ was inactive at doses 25 times greater.

Protein carriers such as lipoproteins enhance targeting of the sensitizer to cancer cells which can overexpress the LDL receptor (Gueddari et al., 1993). Attachment of AlPcS₄ via covalent coupling to the apolipoprotein through a caproic acid spacer chain or via non-covalent adsorption of AlPcS₄(C12) in the phospholipid part of the LDL was used to further determine the utility of LDL as a sensitizer transporter. The AlPcS₄A₂-LDL bioconjugate was only slightly phototoxic *in vitro* with LD₉₀ values of 27 and 33 J cm^{-2} against the EMT-6 and A549 cell lines, respectively. LD₉₀ values of 2 and 4 J cm^{-2} were measured for the non-covalently labeled AlPcS₄(C12)-LDL conjugate using identical experimental conditions. Clearly, LDL-association with lipophilic phthalocyanines via adsorption into its phospholipid bilayer augments the *in vitro* phototoxicity of the Pc. However, *in vivo*, the EMT-6 tumor response of the free AlPcS₄(C12) is similar to that of the LDL-associated dye, most likely due to the known redistribution of amphiphilic photosensitizers to serum lipoproteins upon injection.

Adenovirus proteins were studied as potential Pc carriers. AlPcS₄ was covalently coupled to various adenovirus type 2 capsid proteins, including the hexon, penton base and fiber, via one or two caproic acid spacer chains in 7:1 to 66:1 molar ratios. Adenoviruses gain access to cells via receptor-mediated endocytosis. Adenoviruses

attach to cells via the fiber capsid protein/receptor interaction while cell internalization involves binding of the virus to α_v integrins, a process mediated by RGD sequences found in the penton base of the adenoviral capsid. Cell adsorption studies were carried out on the integrin receptor positive A549 cell line. All Pc bioconjugates exhibited significantly improved cell uptake as compared to free Pc derivatives. In the meanwhile, the photocytotoxicity of the bioconjugates was determined against integrin receptor positive A549 and HEP2 and receptor negative EMT-6 cells. The Pc bioconjugates showed improved activity compared to the parent compounds in the receptor positive cell lines. Pcs labeled with the RGD-containing penton base were most phototoxic requiring roughly half the light dose to induce 50% cell kill as compared to the unlabeled compound. Complete tumor regression of EMT-6 tumors implanted on Balb-c mice was obtained using a mixture of adenovirus type 2 soluble proteins covalently labeled with AlPcS_4A_2 at a dose of $0.5 \mu\text{mol kg}^{-1}$ and 400 J cm^{-2} . Unlabeled AlPcS_4A_2 requires $1 \mu\text{mol kg}^{-1}$ at the same light dose. This suggests that the high affinity RGD/receptor complex is able to target phthalocyanines for PDT.

The effect of an RGD containing peptide on the photodynamic action of $\text{AlPcS}_{2\text{adj}}$ was studied *in vitro* using both a receptor negative and a receptor positive cell line, EMT-6 and A549 cells, respectively. At elevated RGD concentrations, $\geq 10 \text{ mM}$, RGD induced apoptosis. However at lower concentrations, the RGD abrogated the apoptotic effect of PDT in both cell lines, the effect being more pronounced in the A549 cells. At an RGD concentration of $50 \mu\text{M}$, there was over 75 % A549 cell survival following $\text{AlPcS}_{2\text{adj}}$ PDT. Only 30 % survival was observed when no RGD peptide was administered. The protection is seemingly independent of the integrin receptor because it is observed in the

receptor negative cells and the concentrations required are higher than those needed to saturate the receptor.

Further studies are warranted to elucidate the signaling properties of the RGD peptide. Only when we fully understand its role in apoptosis can we strategize to improve current PDT protocols. However, this work has clearly helped phthalocyanines along the road towards the fulfillment of their immense promise as photosensitizers for PDT.

Chapter 7.

Acknowledgements

I wish to extend a sincere thank you to Dr. J. E. van Lier for having the patience to allow me to explore several different pathways. I take with me fond memories and an invaluable education. I would also like to thank all the members of his laboratory who have helped make this thesis possible. I especially thank the chemists for supplying me with ample supplies of various photosensitizers in order to complete the biological assays.

I thank my co-director, Dr. J. M. Weber for his advice and expertise within the adenovirus world.

A heartfelt thank-you to Lise Imbeault, whom without which I never would have made it through. Her technical assistance was greatly appreciated. I will cherish her friendship always. Merci Lise pour ta grande amitié, ta générosité et tes encouragements.

Many thanks to Carole La Madeleine, Stéphanie Marengo and Angèlique Ruzindana for their friendship so dear to me.

Thank you to my mother for her patience, love, strength and support. I hope I have made both my mother and father proud of me. Thanks to my brother Brent for all the encouragement and computer technical assistance.

A very special thank-you to Wesley Sharman who has stood by me through very trying times. His encouragement and strength was invaluable. It was only through his faith in me that I had faith in myself. I am forever indebted to him. I only hope my love is enough. Wesley, Tally and Sammy, you make my world a better place.

I am grateful to the Medical Research Council of Canada for providing the financial support.

Lastly, I wish to thank Dr. B. Paquette, Dr. D. J. Hunting, Dr. R. H. Pottier, Dr. J. M. Weber and Dr. J. E. van Lier for accepting to evaluate this thesis.

Merci à tout le monde qui m'a encouragé et aidé pendant mes études ici à l'Université de Sherbrooke. Merci à tous mes merveilleux amis.

Chapter 8.

Bibliography

- Agarwal, R., M. Athar, D. R. Bickers and H. Mukhtar (1990) Evidence for the involvement of singlet oxygen in the photodestruction by chloroaluminum phthalocyanine tetrasulfonate. *Biochem. Biophys. Res. Commun.* **173**, 34-41.
- Agarwal, R., S. I. Zaidi, M. Athar, D. R. Bickers and H. Mukhtar (1992) Photodynamic effects of chloroaluminum phthalocyanine tetrasulfonate are mediated by singlet oxygen: In vivo and in vitro studies utilizing hepatic microsomes as a model membrane source. *Arch. Biochem. Biophys.* **294**, 30-37.
- Agarwal, R., H. E. Larkin, S. I. A. Zaidi, H. Mukhtar and N. L. Oleinick (1993) Phospholipase activation triggers apoptosis in photosensitized mouse lymphoma cells. *Cancer Res.* **53**, 5897-5902.
- Agarwal, R., N. J. Korman, R. R. Mohan, D. K. Feyes, S. Jawed, M. T. Zaim and H. Mukhtar (1996) Apoptosis is an early event during phthalocyanine photodynamic therapy-induced ablation of chemically induced squamous papillomas in mouse skin. *Photochem. Photobiol.* **63**, 547-552.
- Ahmad, N., D. K. Feyes, R. Agarwal and H. Mukhtar (1998) Photodynamic therapy results in induction of WAF1/CIP1/p21 leading to cell cycle arrest and apoptosis. *Proc. Natl. Acad. Sci. USA* **95**, 6977-6982.
- Ahmad, N., S. Gupta and H. Mukhtar (1999) Involvement of retinoblastoma (Rb) and E2F transcription factors during photodynamic therapy of human epidermoid carcinoma cells A431. *Oncogene* **18**, 1891-1896.
- Ahmad, N. and H. Mukhtar (2000) Mechanism of photodynamic therapy-induced cell death. *Meth. In Enzymol.* **319**, 342-358.
- Akhlynina, T. V., A. A. Rosenkranz, D. A. Jans, P. V. Gulak, N. V. Serebryakova and A. S. Sobolev (1993) The use of internalizable derivatives of chlorin E₆ for increasing its photosensitizing activity. *Photochem. Photobiol.* **58**, 44-48.
- Akhlynina, T. V., A. A. Rosenkranz, D. A. Jans and A. S. Sobolev (1995) Insulin-mediated intracellular targeting enhances the photodynamic activity of chlorin e₆. *Cancer Res.* **55**, 1014-1019.
- Akhlynina, T. V., D. A. Jans, A. A. Rosenkranz, N. V. Statsyuk, I. Y. Balashova, G. Toth, I. Pavo, A. B. Rubin and A. S. Sobolev (1997) Nuclear Targeting of chlorin e₆ enhances its photosensitizing activity. *J. Biol. Chem.* **272**, 20328-20331.
- Akhlynina, T. V., D. A. Jans, N. V. Statsyuk, I. Y. Balashova, G. Toth, I. Pavo, A. A. Rosenkranz, B. S. Naroditshy and A. S. Sobolev (1999) Adenoviruses synergize with nuclear localization signals to enhance nuclear delivery and photodynamic action of internalizable conjugates containing chlorin e₆. *Int. J. Cancer* **81**, 734-740.
- Ali, H., R. Langlois, J. R. Wagner, N. Brasseur, B. Paquette and J. E. van Lier (1988) Biological activities of phthalocyanines X. Syntheses and analyses of sulfonated phthalocyanines. *Photochem. Photobiol.* **47**, 713-717.
- Ali, H. and J. E. van Lier (1999) Metal complexes as photo- and radiosensitizers. *Chem. Rev.* **99**, 2379-2450.
- Alléman, E., R. Gurny and E. Doelker (1993) Drug-loaded nanoparticles – Preparation methods and drug targeting issue. *Eur. J. Pharm. Biopharm.* **39**, 173-191.
- Allémann, S., Kudrevich, K. Lewis, R. Gurny, J. E. van Lier (1995) In vivo photodynamic activity of hexadecafluoro zinc phthalocyanine loaded in peg-coated nanoparticles. *Proc. 1st World Meeting APGI/APV*, Budapest, 9/11 May 1995.

- Alléman, E., J. Rousseau, N. Brasseur, S. V. Kudrevich, K. Lewis and J. E. van Lier (1996) Photodynamic therapy of tumours with hexadecafluoro zinc phthalocyanines formulated in PEG-coated poly(lactic acid) nanoparticles. *Int. J. Cancer* **66**, 821-824.
- Alléman, E., N. Brasseur, O. Benrezzak, J. Rousseau, S. V. Kudrevich, R. W. Boyle, J.-C. Leroux, R. Gurny and J. E. van Lier (1996) PEG-coated poly(lactic acid) nanoparticles for the delivery of hexadecafluoro zinc phthalocyanine to EMT-6 mouse mammary tumours. *J. Pharm. Pharmacol.* **47**, 382-387.
- Alléman, E., N. Brasseur, S. V. Kudrevich, C. La Madeleine and J. E. van Lier (1997) Photodynamic activities and biodistribution of fluorinated zinc phthalocyanines derivatives in the murine EMT-6 tumour model. *Int. J. Cancer* **72**, 289-294.
- Allen, C. M., J. M. Weber and J. E. van Lier (1995) Sulfophthalocyanines for the photodynamic inactivation of viruses in blood products. Effect of structural modifications. *Photochem. Photobiol.* **62**, 184-189.
- Allen, C. M., W. M. Sharman, C. La Madeleine, J. M. Weber, R. Langlois, R. Ouellet and J. E. van Lier (1999) Photodynamic therapy: tumor targeting with adenoviral proteins. *Photochem. Photobiol.* **70**, 512-523.
- Allen, C. M., W. M. Sharman and J. E. van Lier (2001) Current status of phthalocyanines in the photodynamic therapy of cancer. *J. Porphyrins Phthalocyanines* **5**, 161-169.
- Allison, B. A., E. Waterfield, A. M. Richter and J. G. Levy (1991) The effects of plasma lipoproteins on *in vitro* tumor cell killing and *in vivo* tumor photosensitization with benzoporphyrin derivative. *Photochem. Photobiol.* **54**, 709-715.
- Allison, B. A., P. H. Pritchard and J. G. Levy (1994) Evidence for low-density lipoprotein receptor-mediated uptake of benzoporphyrin derivative. *Br. J. Cancer* **69**, 833-839.
- Anderson, C. Y., S. Hrabovsky, Y. McKinley, K. Tubesing, H.-P. Tang, R. Dunbar, H. Mukhtar and C. A. Elmetts (1997) Phthalocyanine photodynamic therapy: Disparate effects of pharmacologic inhibitors on cutaneous photosensitivity and on tumor regression. *Photochem. Photobiol.* **65**, 895-901.
- Anderson, C. Y., K. Freye, K. A. Tubesing, Y.-S. Li, M. E. Kenney, H. Mukhtar and C. Elmetts (1998) A comparative analysis of silicon phthalocyanine photosensitizers for *in vivo* photodynamic therapy of RIF-1 tumors in C3H mice. *Photochem. Photobiol.* **67**, 332-336.
- Anderson, V. C. and D. H. Thompson (1992) Triggered release of hydrophilic agents from plasmalogen liposomes using visible light or acid. *Biochimica Biophysica Acta* **1109**, 33-42.
- Arap, W., R. Pasqualini and E. Ruoslahti (1998) Cancer treatment by targeted drug delivery to tumor vasculature in a mouse model. *Science* **279**, 377-380.
- Babincová, M., O. Sourivong and P. Babinec (2000) Gene transfer-mediated intracellular photodynamic therapy. *Med. Hypothesis* **54**, 180-181.
- Bachor, R., C. R. Shea, R. Gillies and T. Hasan (1991) Photosensitized destruction of human bladder carcinoma cells treated with chlorin e_6 -conjugated microspheres. *Proc. Natl. Acad. Sci. USA* **88**, 1580-1584.
- Bachor, R., C. R. Shea, S. J. Belmonte and T. Hasan (1991) Free and conjugated chlorin E_6 in the photodynamic therapy of human bladder carcinoma cells. *J. Urol.* **146**, 1654-1658.
- Bachor, R., M. Scholz, C. R. Shea and T. Hasan (1991) Mechanism of photosensitization by microsphere-bound chlorin e_6 in human bladder carcinoma cells. *Cancer Res.* **51**, 4410-4414.
- Bagnarello, A.G., L. A. Lewis, M. C. McHenry, A. J. Weinstein, H. K. Naito, A. J. McCullough, R. J. Lederman and T. L. Gavan (1977) Unusual serum lipoprotein abnormality induced by the vehicle of miconazole. *N. Engl. J. Med.* **296**, 497-499.

- Bai, M., B. Harfe and P. Freimuth (1993) Mutations that alter an Arg-Gly-Asp (RGD) sequence in the adenovirus type 2 penton base protein abolish its cell-rounding activity and delay virus reproduction in flat cells. *J. Virol.* **67**, 5198-5205.
- Bai, M., L. Campisi and P. Freimuth (1994) Vitronectin receptor antibodies inhibit infection of HeLa and A549 cells by adenovirus type 12 but not by adenovirus type 2. *J. Virol.* **68**, 5925-5932.
- Baker, A. and J. R. Kanofsky (1992) Quenching of singlet oxygen by biomolecules from L1210 leukemia cells. *Photochem. Photobiol.* **55**, 523-528.
- Barinaga, M. (1998) Peptide-guided cancer drugs show promise in mice. *Science* **279**, 323-324.
- Barrett, P. A., C. E. Dent, R. P. Linstead (1936) Phthalocyanines. Part VII. Phthalocyanines as a co-ordinating group. A general investigation of the metallic derivatives. *J. Chem. Soc.*, 1719-1736.
- Barrett, P. A., D. A. Frye, R. P. Linstead (1938) Phthalocyanines and associated compounds. Part XIV. Further investigations of metallic derivatives. *J. Chem. Soc.*, 1157-1163.
- Bates, R. C., L. F. Lincz and G. F. Burns (1995) Involvement of integrins in cell survival. *Cancer and Metastasis Reviews* **14**, 191-203.
- Belin, M.-T. and P. Boulanger (1993) Involvement of cellular adhesion sequences in the attachment of adenovirus to the HeLa cell surface. *J. Gen. Virol.* **74**, 1485-1497.
- Ben Hur, E., J. A. Siwecki, H. C. Newman, S. W. Crane and I. Rosenthal (1987) Mechanism of uptake of sulfonated metallophthalocyanines by cultured mammalian cells. *Cancer Lett.* **38**, 215-222.
- Benfield, J. R., W. G. Hammond, R. R. Paladuga, H. Y. Pak, N. Azumi and R. L. Teplitz (1986) Endobronchial carcinogenesis in dogs. *J. Thoracic Cardiovasc. Surg.* **92**, 880-889.
- Berg, K., J. C. Bommer and J. Moan (1989) Evaluation of sulfonated aluminum phthalocyanines for use in photochemotherapy. A study on the relative efficiencies of photoinactivation. *Photochem. Photobiol.* **49**, 587-594.
- Berg, K., J. C. Bommer and J. Moan (1989) Evaluation of sulphonated aluminum phthalocyanines for use in photochemotherapy. Cellular uptake studies. *Cancer Lett.* **44**, 7-15.
- Berg, K., P. K. Selbo, L. Prasmickaite, T. T. Tjelle, K. Sandvig, J. Moan, G. Gaudernack, Ø. Fodstad, S. Kjølrsrud, H. Anholt, G. H. Rodal, S. K. Rodal and A. Høgset (1999) Photochemical internalization: A novel technology for delivery of macromolecules into cytosol. *Cancer Res.* **59**, 1180-1183.
- Berg, O. G., B.-Z. Yu, J. Rogers and M. K. Jain (1991) Interfacial catalysis by phospholipase A₂: Determination of the interfacial kinetic rate constant. *Biochem.* **30**, 7283-7297.
- Bergelson, J. M., J. A. Cunningham, G. Droguett, E. A. Kurt-Jones, A. Krithivas, J. S. Hong, M. S. Horwitz, R. L. Crowell and R. W. Finberg (1997) Isolation of a common receptor for Coxsackie B virus and adenovirus 2 and 5. *Science* **275**, 1320-1323.
- Bertoloni, G., R. Rossi, G. Valduga, G. Jor, H. Ali, J. E. van Lier (1992) Photosensitizing activity of water- and lipid-soluble phthalocyanines on prokaryotic and eukaryotic microbial cells. *Microbios.* **71**, 33-46.
- Bible, K. C. and S. H. Kaufmann (1996) Flavopiridol: A cytotoxic flavone that induces cell death in noncycling A549 human lung carcinoma cells. *Cancer Res.* **56**, 4856-4861.
- Billetta, R., P. Lanza and M. Zanetti (1997) Ligand function of antigenized antibodies expressing the RGD motif. *Chem. Immunol.* **65**, 159-178.
- Bisland, S. K., D. Singh and J. Garipey (1999) Potentiation of chlorin e6 photodynamic activity in vitro with peptide-based intracellular vehicles. *Bioconjugate Chem.* **10**, 982-992.
- Bissonette, R. and H. Lui (1997) Current status of photodynamic therapy in dermatology. *Lasers in Dermatol.* **15**, 507-519.
- Bogden, A. E., H. J. Esber, D. J. Taylor and J. H. Gray (1974) Comparative study on the effects of surgery, chemotherapy, and immunotherapy, alone and in combination, on metastases of 13762 mammary adenocarcinoma. *Cancer Res.* **34**, 1627-1631.

- Bonnett, R. (1995) Photosensitizers of the porphyrin and phthalocyanine series for photodynamic therapy. *Chem. Soc. Rev.* **24**, 19-33.
- Bonnett, R. (1999) Photodynamic therapy in historical perspective. *Rev. Contemp. Pharmacother.* **10**, 1-17.
- Boulanger, P. A. and F. Puvion (1973) Large-scale preparation of soluble adenovirus hexon, penton and fiber antigens in highly purified form. *Eur. J. Biochem.* **39**, 37-42.
- Bourdon, O., V. Mosqueira, Ph. Legrand and J. Blais (2000) A comparative study of the cellular uptake, localization and phototoxicity of *meta*-tetra(hydroxyphenyl)chlorin encapsulated in surface-modified submicronic oil/water carriers in HT29 tumor cells. *J Photochem. Photobiol. B: Biol.* **55**, 164-171.
- Bowser, M. T., E. D. Sternberg and D. D. Y. Chen (1996) Development and application of a nonaqueous capillary electrophoresis system for the analysis of porphyrins and their oligomers. *Analyt. Biochem.* **241**, 143-150.
- Boyle, RW and D Dolphin (1996) Structure and biodistribution relationships of photodynamic sensitizers. *Photochem. Photobiol.* **63**, 469-485.
- Bozzo, C., G. Bellomo, L. Silengo, G. Tarone and F. Altruda (1997) Soluble integrin ligands and growth factors independently rescue neuroblastoma cells from apoptosis under nonadherent conditions. *Experimental Cell Res.* **237**, 326-337.
- Bradford, M. M. (1976) A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**, 248-254.
- Brasseur, N., H. Ali, R. Langlois, J. R. Wagner, J. Rousseau, J. E. van Lier (1987) Biological activities of phthalocyanines-V. Photodynamic therapy of EMT-6 mammary tumors in mice with sulfonated phthalocyanines. *Photochem. Photobiol.* **45**, 581-586.
- Brasseur, N., H. Ali, R. Langlois and J. E. van Lier (1987) Biological activities of phthalocyanines VII. Photoinactivation of V-79 Chinese hamster cells by selectively sulfonated gallium phthalocyanines. *Photochem. Photobiol.* **46**, 739-744.
- Brasseur, N., H. Ali, R. Langlois and J. E. van Lier (1988) Biological activities of phthalocyanines IX. Photosensitization of V-79 Chinese hamster cells and EMT-6 mouse mammary tumors by selectively sulfonated zinc phthalocyanines. *Photochem. Photobiol.* **47**, 705-711.
- Brasseur, N., D. Brault and D. Couvreur (1991) Adsorption of hematoporphyrin onto polyalkylcyanoacrylate nanoparticles: Carrier capacity and drug release. *Int. J. Pharm.* **70**, 129-135.
- Brasseur, N., T. L. Nguyen, R. Langlois, R. Ouellet, S. Marengo, D. Houde, J. E. van Lier (1994) Synthesis and photodynamic activities of silicon 2,3-naphthalocyanine derivatives. *J. Med. Chem.* **37**, 415-420.
- Brasseur, N., R. Ouellet, K. Lewis, W. R. Potter, J. E. van Lier (1995) Photodynamic activities and skin photosensitivity of *bis*(dimethylhexylsiloxy)silicon 2,3-naphthalocyanine in mice. *Photochem. Photobiol.* **62**, 1058-1065.
- Brasseur, N., R. Langlois, C. La Madeleine, R. Ouellet and J. E. van Lier (1999) Receptor-mediated targeting of phthalocyanines to macrophages via covalent coupling to native or maleylated bovine serum albumin. *Photochem. Photobiol.* **63**, 345-352.
- Brasseur, N., R. Ouellet, C. La Madeleine and J. E. van Lier (1999) Water-soluble aluminium phthalocyanine-polymer conjugates for PDT: Photodynamic activity and pharmacokinetics in tumour-bearing mice. *Br. J. Cancer* **80**, 1533-1541.
- Braun, A. and J. Tscherniac (1907) Products of the action of acetic anhydride on phthalamide. *Ber.* **40**, 2709-2714.
- Broers, J. L. V. and F. C. S. Ramaekes (1994) Differentiation markers for lung-cancer sub-types. A comparative study of their expression *in vivo* and *in vitro*. *Int. J. Cancer* **S8**, 134-137.

- Brooks, P. C., A. M. P. Montgomery, M. Rosenfeld, R. A. Resifeld, T. Hu, G. Klier and D. A. Cheresh (1994) Integrin $\alpha v \beta 3$ antagonists promote tumor regression by inducing apoptosis of angiogenic blood vessels. *Cell* **79**, 1157-1164.
- Brooks, P. C., R. A. F. Clark and D. A. Cheresh (1994) Requirement of vascular integrin $\alpha v \beta 3$ for angiogenesis. *Science* **264**, 569-571.
- Brooks, P. C., S. Silletti, T. L. von Schalscha, M. Friedlander and D. A. Cheresh (1998) Disruption of angiogenesis by PEX, a noncatalytic metalloproteinase fragment with integrin binding activity. *Cell* **92**, 391-400.
- Brown, J. E., S. Brown and D. I. Vernon (1999) Photodynamic therapy- New light on cancer treatment. *JSDC* **115**, 249-253.
- Brown, S. B. and T. G. Truscott (1993) New light on cancer therapy. *Chem. in Br.* **29**, 955-958.
- Bru, R., J. M. López-Nicolás, E. Núñez-Delicado, D. Nortes-Ruipérez, A. Sánchez-Ferrer and F. García-Carmona (1997) Cyclodextrins as hosts for poorly water-soluble compounds in enzyme catalysis. *Appl. Biochem. Biotech.* **61**, 189-198.
- Buckley, C. D., D. Pilling, N. V. Henriquez, G. Parsonage, K. Threlfall, D. Scheel-Toellner, D. L. Simmons, A. N. Akbar, J. M. Lord and M. Salmon (1999) RGD peptides induce apoptosis by direct caspase-3 activation. *Nature* **397**, 534-539.
- Buettner, G. R. and B. A. Jurkiewicz (1996) Catalytic metals, ascorbate and free radicals: Combinations to avoid. *Rad. Res.* **145**, 532-541.
- Bylund, D. B. and H. I. Yamamura (1990) Methods for receptor binding. In: *Methods in Neurotransmitter receptor analysis*. Eds.: H. I. Yamamura et. al., New York: Raven Press Ltd., p.1-35.
- Bylund, D. B. and M. L. Toews (1993) Radioligand binding methods: practical guide and tips. *Am. J. Physiol.* **265**, L421-L429.
- Byrne, G. T., R. P. Linstead, A. R. Lowe (1934) Phthalocyanines. Part II. The preparation of phthalocyanines and some metallic derivatives from o-cyanobenzamide and phthalimide. *J. Chem. Soc.* 1017-1022.
- Calvete, J. J. (1994) Clues for understanding the structure and function of a prototypic human integrin: The platelet glycoprotein IIb/IIIa complex. *Thrombosis and Haemostasis* **72**, 1-15.
- Canon, J. B. (1993) Pharmaceuticals and drug delivery aspects of heme and porphyrin therapy. *J. Pharmaceut. Sci.* **82**, 435-446.
- Canti, G., P. Franco, O. Marelli, R. Cubeddu, P. Taroni and R. Ramponi (1990) Comparative study of the therapeutic effect of photoactivated hematoporphyrin derivative and aluminium disulfonated phthalocyanines on tumor bearing mice. *Cancer Lett.* **53**, 123-127.
- Carcenac, M., C. Larroque, R. Langlois, J. E. van Lier, J.-C. Artus and A. Pèlerin (1999) Preparation, phototoxicity and biodistribution studies of anti-carcinoembryonic antigen monoclonal antibody-phthalocyanine conjugates. *Photochem. Photobiol.* **70**, 930-936.
- Cardone, M. H., N. Roy, H. R. Stennicke, G. S. Salvesen, T. F. Franke, E. Stanbridge, S. Frisch and J. C. Reed (1998) Regulation of cell death protease caspase-9 by phosphorylation. *Science* **282**, 1318-1321.
- Carmichael, J., W. G. DeGraff, A. F. Gazdar, J. D. Minna and J. B. Mitchell (1987) Evaluation of a tetrazolium-based semi-automated colorimetric assay: assessment of radiosensitivity. *Cancer Res.* **47**, 943-946.
- Carpenter, S., M. J. Fehr, G. A. Kraus and J. W. Petrich (1994) Chemiluminescent activation of the activiral activity of hypericin: a molecular flashlight. *Proc. Natl. Acad. Sci. USA* **91**, 12273-12277.
- Chan, W.-H., J.-S. Yu and S.-D. Yang (2000) Apoptotic signalling cascade in photosensitized human epidermal carcinoma A431 cells: Involvement of singlet oxygen, c-Jun N-terminal kinase, caspase-3 and p21-mediated kinase 2. *Biochem. J.* **351**, 221-232.

- Chan, W.-S., J. F. Marshall, R. Svensen, D. Phillips and I. R. Hart (1987) Photosensitizing activity of phthalocyanine dyes screened against tissue culture cells. *Photochem. Photobiol.* **45**, 757-761.
- Chan, W.-S., C. M. L. West, J. V. Moore and I. R. Hart (1991) Photocytotoxic efficacy of sulphonated species of aluminium phthalocyanine against cell monolayers, multicellular spheroids and *in vivo* tumours. *Br. J. Cancer* **64**, 827-832.
- Chan, W.-S., N. Brasseur, C. La Madeleine and J. E. van Lier (1997) Evidence for different mechanisms of EMT-6 tumor necrosis by photodynamic therapy with disulfonated aluminum phthalocyanine or photofrin: tumor cell survival and blood flow. *Anticancer Res.* **16**, 1887-1892.
- Chan, W.-S., N. Brasseur, C. La Madeleine, R. Ouellet and J. E. van Lier (1997) Efficacy and mechanism of aluminium phthalocyanine and its sulphonated derivatives mediated photodynamic therapy on murine tumours. *Eur. J. Cancer* **33**, 1855-1859.
- Chapman, H. A. (1997) Plasminogen activators, integrins, and the coordinated regulation of cell adhesion and migration. *Curr. Opin. Cell Biol.* **9**, 714-724.
- Chen, C., S. Parangi, M. J. Tolentino and J. Folkman (1995) A strategy to discover circulation angiogenesis inhibitors generated by human tumors. *Cancer Res.* **55**, 4230-4233.
- Choi Kim, H.-R., Y. Luo, G. Li and D. Kessel (1999) Enhanced apoptotic response to photodynamic therapy after bcl-2 transfection. *Cancer Res.* **59**, 3429-3432.
- Clarke, A. S., M. M. Lotz, C. Chao, and A. M. Mercurio (1995) Activation of the p21 pathway of growth arrest and apoptosis by the β_4 integrin cytoplasmic domain. *J. Biol. Chem.* **270**, 22673-22676.
- Conforti, G., C. Dominguez-Jimenez, A. Zanetti, M. A. Gimbrone Jr., O. Cremona, P. C. Marchisio and E. Dejana (1992) Human endothelial cells express integrin receptors on the luminal aspect of their membrane. *Blood* **80**, 437-446.
- Cook, M. J., I. Chambrier, S. J. Cracknell, D. A. Mayes and D. A. Russell (1995) Octa-alkyl zinc phthalocyanines: Potential photosensitizers for use in the photodynamic therapy of cancer. *Photochem. Photobiol.* **62**, 542-545.
- Cotton, M. and J. M. Weber (1995) The adenovirus protease is required for virus entry into host cells. *Virology* **213**, 494-502.
- Curiel, D. T., S. Agarwal, E. Wagner and M. Cotton (1991) Adenovirus enhancement of transferrin-polylysine mediated gene delivery. *Proc. Natl. Acad. Sci. USA* **88**, 8850-8854.
- Cuzange, A., J. Chroboczek and B. Jacrot (1994) The penton base of human adenovirus type 3 has the RGD motif. *Gene* **146**, 257-259.
- Dahle, J., H. B. Steen and J. Moan (1999) The mode of cell death induced by photodynamic treatment depends on cell density. *Photochem. Photobiol.* **70**, 363-367.
- Damiano, J. S., A. E. Cress, L. A. Hazlehurst, A. A. Shtil and W. S. Dalton (1999) Cell adhesion mediated drug resistance (CAM-DR): Role of integrins and resistance to apoptosis in human myeloma cell lines. *Blood* **39**, 1658-1667.
- Dandridge, A. G., H. A. Drescher, J. Thomas (1929) Dyes. *British Patent* 322,169 (Scottish Dyes, Ltd.), May 16.
- Danen, E. H. J., K. F. J. Jansen, A. A. Van Kraats, I. M. H. A. Cornelissen, D. J. Ruiter and G. N. P. Van Muijen (1995) α_v -Integrins in human melanoma: gain of $\alpha_v\beta_3$ and loss of $\alpha_v\beta_5$ are related to tumor progression *in situ* but not to metastatic capacity of cell lines in nude mice. *Int. J. Cancer* **61**, 491-496.
- Daniell, M. D. and J. S. Hill (1991) A history of photodynamic activity. *Aust. N. Z. J. Surg.* **61**, 340-348.
- Darwent, J. R., P. Douglas, A. Harriman, G. Porter and M.-C. Richoux (1982) Metal phthalocyanines and porphyrins as photosensitizers for reduction of water to hydrogen. *Coord. Chem. Rev.* **44**, 83-126.
- Davies, J. S., C. Enjalbal, C. Wise, G. Jones and B. K. Handa (1994) Further studies on cyclic RGD analogues. *Biochem. Soc. Trans.* **22**, 1049-1052.

- Davis, N., D. Liu, A. K. Jain, S. Y. Jiang, A. Richter and J. Levy (1993) Modified polyvinyl alcohol-benzoporphyrin derivative conjugates as phototoxic agents. *Photochem. Photobiol.* **57**, 641-647.
- Daziano, J.-P., S. Steenken, C. Chabannon, P. Mannoni, M. Chanon and M. Julliard. (1996) Photophysical and redox properties of a series of phthalocyanines: Relation with their photodynamic activities on *TF-1* and Daudi Leukemic cells. *Photochem. Photobiol.* **64**, 712-719.
- de Diesbach, H. and E. von der Weid (1927) Quelques sels complexes des o-dinitriles avec le cuivre et la pyridine. *Helv. Chim. Acta* **10**, 886-888.
- de Smidt, P. C., A. J. Versluis and T. J. C. van Berkel (1993) Properties of incorporation, redistribution, and integrity of porphyrin-low-density lipoprotein complexes. *Biochem.* **32**, 2916-2922.
- de Vries, H. E., A. C. E. Moor, T. M. A. R. Dubbelman, T. J. C. van Berkel and J. Kuiper (1999) Oxidized low-density lipoprotein as a delivery system for photosensitizers: Implications for photodynamic therapy of atherosclerosis. *J. Pharmacol. Exp. Therapeut.* **289**, 528-534.
- Decreau, R., M. J. Richard, P. Verrando, M. Chanon and M. Juilliard (1999) Photodynamic activities of silicon phthalocyanine against achromic M6 melanoma cells and healthy human melanocytes and keratinocytes. *J. Photochem. Photobiol. B Biol.* **48**, 48-56.
- Dedhar, S. and G. E. Hannigan (1996) Integrin cytoplasmic interactions and bidirectional transmembrane signalling. *Curr. Opinion Cell Biol.* **8**, 657-669.
- del Governatore, M., M. R. Hamblin, C. R. Shea, I. Rizvi, K. G. Molpus, K. K. Tanabe and T. Hasan (2000) Experimental photoimmunotherapy of hepatic metastases of colorectal cancer with a 17.1A chlorin_{e6} immunoconjugates. *Cancer Res.* **60**, 4200-4205.
- del Governatore, M., M. R. Hamblin, E. E. Piccinini, G. Ugolini and T. Hasan (2000) Targeted photodestruction of human colon cancer cells using charged 17.1A chlorin *e*₆ immunoconjugates. *Br. J. Cancer* **82**, 56-64.
- Dellinger, M. (1996) Apoptosis or necrosis following Photofrin photosensitization: influence of the incubation protocol. *Photochem. Photobiol.* **64**, 182-187.
- Dent, C. E. and R. P. Linstead (1934) Phthalocyanines. Part IV. Copper phthalocyanines. *J. Chem. Soc.* 1027-1031.
- Dent, C. E., R. P. Linstead and A. R. Lowe (1934) Phthalocyanines. Part VI. The structure of the phthalocyanines. *J. Chem. Soc.* 1033-1039.
- Dent, C. E. (1938) Preparation of phthalocyanine-like pigments related to the porphyrins. *J. Chem. Soc.* 1-6.
- Dereski, M. O., L. Madigan and M. Chopp (1995) Brain response to photodynamic therapy with Photofrin, nonsulfonated aluminum phthalocyanine and tin purpurin. *Proc. SPIE* **2371**, 579-581.
- Devanathan, S., T. A. Dahl, W. R. Midden and D. C. Neckers (1990) Readily available fluorescein isothiocyanate-conjugated antibodies can be easily converted into targeted phototoxic agents for antibacterial, antiviral, and anticancer therapy. *Proc. Natl. Acad. Sci. USA* **87**, 2980-2984.
- Donati, G., A. D. Kapetanios and C. J. Pournaras (1999) Principles of treatment of choroidal neovascularization with photodynamic therapy in age-related macular degeneration. *Sem. Ophthalmol.* **14**, 2-10.
- Dorr, R. T. (1994) Pharmacology and toxicology of cremophor EL diluent. *Ann. Pharmacotherapy* **28**, S11-S14.
- Dougherty, T. J., S. G. Grindley, R. Fiel, K. R. Weishaupt and D. G. Boyle (1975) Photoradiation therapy II. Cure of animal tumours with hematoporphyrin and light. *J. Natl. Cancer Inst.* **55**, 115-119.
- Dougherty, T. J., J. E. Kaufman, A. Goldfarb, K. R. Weishaupt and D. G. Boyle (1978) Photoradiation therapy for the treatment of malignant tumours. *Cancer Res.* **38**, 2628-2635.

- Dougherty, T. J., C. J. Gomer, B. W. Henderson, G. Jori, D. Kessel, M. Korblick, J. Moan and Q. Peng (1998) Photodynamic therapy. *J. Natl. Can. Inst.* **90**, 889-905.
- Doumas, B. T., W. Watson and H. G. Biggs (1971) Albumin standards and the measurement of serum albumin with bromocresol green *Clinica Chimica Acta*. **31**, 97-107.
- Dummin, H., Th. Cernay and H. W. Zimmermann (1997) Selective photosensitization of mitochondria in HeLa cells by cationic Zn(II) phthalocyanines with lipophilic side-chains. *J. Photochem. Photobiol. B: Biol.* **37**, 219-229.
- Duska, L. R., M. R. Hamblin, M. P. Bamberg and T. Hasan (1997) Biodistribution of charged F(ab')₂ photoimmunoconjugates in a xenograft model of ovarian cancer. *Br. J. Cancer* **75**, 837-844.
- Dye, D. and J. Watkins (1980) Suspected anaphylactic reaction to Cremophor EL. *Br. Med. J.* **280**, 1353.
- Earnshaw, W. C., L. M. Martins and S. H. Kaufmann (1999) Mammalian caspases: Structure, activation, substrates and functions during apoptosis. *Annu. Rev. Biochem.* **68**, 383-424.
- Edrei, R., V. Gottfried, J. E. van Lier and S. Kimel (1998) Sulfonated phthalocyanines: Photophysical properties, *in vitro* cell uptake and structure-activity relationships. *J. Porphyrins Phthalocyanines* **2**, 191-199.
- Egorin, M. J., E. G. Zuhowski, D. L. Sentz, J. M. Dobson, P. S. Callery and J. L. Eiseman (1999) Plasma pharmacokinetics and tissue distribution in CD2F1 mice of Pc4 (NSC 676418), a silicone phthalocyanine photodynamic sensitizing agent. *Cancer Chemother. Pharmacol.* **44**, 283-294.
- Egorov, S. I. U., A. A. Krasnovskii Jr, D. B. Pankovskii, G. V. Ponomarev, and A. P. Savitskii (1990) [Photosensitized formation of singlet oxygen in water solutions of covalent porphyrin-antibody conjugates (Russian)]. *Biulleten Eksperimentalnoi Biologii I Meditsiny* **109**, 349-351.
- Ellerby, H. M., W. Arap, L. M. Ellerby, R. Kain, R. Andrusiak, G. Del Rio, S. Krajewski, C. R. Lombardo, R. Rao, E. Ruoslahti, D. E. Bredesen and R. Pasqualini (1999) Anti-cancer activity of targeted proapoptotic peptides *Nature Med.* **5**, 1032-1038.
- Engleman, V. W., G. A. Nickols, F. P. Ross, M. A. Horton, D. W. Griggs, S. L. Settle, P. G. Ruminiski and S. L. Teitelbaum (1997) A peptidomimetic antagonists of the α sub v β sub 3 integrin inhibits bone resorption *in vitro* and prevents osteoporosis *in vivo*. *J. Clin. Invest.* **99**, 2284-2292.
- Evans, H. H., M.-F. Horng, M. Ricanati, J. T. Deahl, N. L. Oleinick (1997) Mutagenicity of photodynamic therapy as compared to UVC and ionizing radiation in human murine lymphoblast cell lines. *Photochem. Photobiol.* **66**, 690-696.
- Fairbanks, G., T. L. Steck and D. F. Wallach (1971) Electrophoretic analysis of the major polypeptides of the human erythrocyte membrane. *Biochem.* **10**, 2606-2617.
- Falcioni, R., L. Cimino, M. P. Gentileschi, I. D'Agnano, G. Zupi, S. J. Kennel and A. Sacchi (1994) Expression of β 1, β 3, β 4, and β 5 integrins by human lung carcinoma cells of different histologies. *Exp. Cell Res.* **210**, 113-122.
- Faull, R. J., N. L. Kovach, J. M. Harlan and M. H. Ginsberg (1994) Stimulation of integrin-mediated adhesion of T lymphocytes and monocytes: Two mechanisms with divergent biological consequences. *J. Exp. Med.* **179**, 1307-1316.
- Feix, J. B., M. S. Anderson and B. Kalyanaraman (1994) Enhancement of merocyanine 540 uptake and photodynamic cell killing by salicylates. *Cancer Res.* **54**, 3474-3478.
- Felding-Habermann, B., B. M. Mueller, C. A. Romerdahl and D. A. Cheresh (1992) Involvement of integrin α v gene expression in human melanoma tumorigenicity. *J. Clin. Invest.* **89**, 2018-2022.
- Fernandez, J. M., M. D. Bilgin, L. I. Grossweiner (1997) Singlet oxygen generation by photodynamic agents. *J. Photochem. Photobiol. B: Biol.* **37**, 131-140.

- Fingert, H. J., J. Campisi and A. B. Pardee (1997) Cell proliferation and differentiation. In *Cancer Medicine*. (Eds. J. F. Holland, R. C. Bast Jr, D. L. Morton, E. Frei III, D. W. Kufe and R. R. Weichselbaum) William and Wilkins: Baltimore; 4th ed, pp.3-8.
- Fisher, A. M. R., K. Danenberg, D. Banerjee, J. R. Bertino, P. Danenberg and C. J. Gomer (1997) Increased photosensitivity in HL60 cells expressing wild-type p53. *Photochem. Photobiol.* **66**, 265-270.
- Fisher, A. M. R., A. Ferrario, N. Rucker, S. Zhang and C. J. Gomer (1999) Photodynamic therapy sensitivity is not altered in human tumor cells after abrogation of p53 function. *Cancer Res.* **59**, 331-335.
- Flynn, G., T. J. Hackett, L. McHale and A. P. McHale (1994) Magnetically responsive photosensitizing reagents for possible use in photoradiation therapy. *Cancer Lett.* **78**, 109-114.
- Foley, M. S. C., A. Beeby, A. W. Parker, S. M. Bishop and D. Phillips (1997) Photophysics of disulphonated aluminium phthalocyanine in reverse micelles of Aerosol OT. *J. Photochem. Photobiol. B: Biol.* **38**, 18-24.
- Foley, M. S. C., A. Beeby, A. W. Parker, S. M. Bishop and D. Phillips (1997) Excited triplet state photophysics of the sulphonated aluminium phthalocyanines bound to human serum albumin. *J. Photochem. Photobiol. B: Biol.* **38**, 10-17.
- Folkman, J. (1995) Seminars in medicine of the Beth Israel Hospital, Boston. Clinical applications of research on angiogenesis. *New Engl. J. Med.* **333**, 1757-1763.
- Folkman, J. (1996) Fighting Cancer by attacking its blood supply. *Scientific American* **275**, 150-154.
- Folkman, J. (1996) New perspectives in clinical oncology from angiogenesis research. *Eur. J. Cancer* **32A**, 2534-2539.
- Folkman, J. and P. A. D'Amore (1996) Blood vessel formation: What is its molecular basis? *Cell* **87**, 1153-1155.
- Folkman, J. (1997) Addressing tumor blood vessels. *Nature Biotech.* **15**, 510.
- Foote, C. S. (1976) Photosensitized oxidation and singlet oxygen: Consequences in biological systems. In *Free Radicals in Biology* (Edited by W. A. Pryor), Vol. II, pp. 85-133. Academic Press, New York.
- Foote, C. S. (1991) Guest editorial: Definition of Type I and Type II photosensitized oxidation. *Photochem. Photobiol.* **54**, 659.
- Fowler, G. J. S. and R. Devonshire (1992) Photobleaching of 1,3-diphenylisobenzofuran by novel phthalocyanine dye derivative. *J. Photochem. Photobiol. B: Biol.* **14**, 177-185.
- Freimuth, P. (1996) A human cell line selected for resistance to adenovirus infection has reduced levels of the virus receptor. *J. Virol.* **70**, 4081-4085.
- Friedberg, J. S., R. G. Tompkins, S. L. Rakestraw, S. W. Warren, A. J. Fischman and M. L. Yarmush (1991) Antibody-targeted photolysis: Bacteriocidal effects of Sn(IV) chlorin e6-dextran-monoclonal antibody conjugates. *Annal. NY Acad. Sci.* **618**, 383-393.
- Frisch, S. M. and E. Ruoslahti (1997) Integrins and anoikis. *Curr. Opinion Cell Biol.* **9**, 701-706.
- Fu, Y. and J. R. Kanofsky (1995) Singlet oxygen generation from liposomes: A comparison of time-resolved 1270 nm emission with singlet-oxygen kinetics calculated from a one dimensional model of singlet-oxygen diffusion and quenching. *Photochem. Photobiol.* **62**, 692-702.
- Fuchs, J. and J. Thiele (1998) The role of oxygen in cutaneous photodynamic therapy. *Free Rad. Biol. Med.* **24**, 835-847.
- Gagnebin, J., M. Brunori, M. Otter, L. Juillerat-Jeanneret, P. Monnier and R. Iggo (1999) A photosensitising adenovirus for photodynamic therapy. *Gene Therapy* **6**, 1742-1750.
- Gal, D., P. C. MacDonald, J. C. Porter, J. W. Smith and E. R. Simpson (1981) Effect of cell density and confluency on cholesterol metabolism in cancer cells in monolayer culture. *Cancer Res.* **41**, 473-477.

- Gantchev, T. G., I. J. Urumov, D. J. Hunting and J. E. van Lier (1994) Photocytotoxicity and intracellular generation of free radicals by tetrasulphonated Al- and Zn-phthalocyanines. *Int. J. Radiat. Biol.* **65**, 289-298.
- Gantchev, T. G. and J. E. van Lier (1995) Catalase inactivation following photosensitization with tetrasulfonated metallophthalocyanines. *Photochem. Photobiol.* **62**, 123-134.
- Garbo, G. M. (1990) The use of liposomes, emulsions or inclusion complexes may potentiate in vivo effects of SnET2. *Proc. Int. Soc. Optical Eng.* **1203**, 118-125.
- Gaspard, S., P. Margaron, C. Tempête and T. -H. Tran Thi (1990) Mixed acenannellated metallotetraazaporphins: A new class of amphiphilic photosensitizers for photodynamic therapy of cancer. *J. Photochem. Photobiol. B: Biol.* **4**, 419-423.
- Gaspard, S., C. Tempête and G. H. Werner (1995) Studies on photoinactivation by various phthalocyanines of a free or replication non-enveloped virus. *J. Photochem. Photobiol. B: Biol.* **31**, 159-162.
- Giancotti, F. G. (1997) Integrin signaling: specificity and control of cell survival and cell cycle progression. *Curr. Opinion Cell Biol.* **9**, 691-700.
- Giancotti, F. G. and E. Ruoslahti (1999) Integrin Signaling. *Science* **285**, 1028-1032.
- Gijsens, A. and P. De Witte (1998) Photocytotoxic action of EGF-PVA-Sn(IV)chlorine6 and EGF-dextran-Sn(IV)chlorine6 internalizable conjugates on A431 cells. *Int. J. Oncol.* **13**, 1171-1177.
- Gijsens, A., L. Missiaen, W. Merlevede and P. de Witte (2000) Epidermal growth factor-mediated targeting of chlorin e₆ selectively potentiates its photodynamic activity. *Cancer Res.* **60**, 2197-2202.
- Ginevra, F. S. Bifflanti, A. Pagnan, R. Biolo, E. Reddi and G. Jori (1990) Delivery of the tumour photosensitizer zinc(II) phthalocyanine to serum proteins by different liposomes: studies *in vitro* and *in vivo*. *Cancer Lett.* **49**, 59-65.
- Girotti, A. W. (1975) Photodynamic action of bilirubin on human erythrocyte membranes. Modification of polypeptide constituents. *Biochem.* **14**, 3377-3383.
- Goff, B. A., M. P. Bamberg and T. Hasan (1991) Photoimmunotherapy of human ovarian carcinoma cell *ex vivo*. *Cancer Res.* **51**, 4762-4767.
- Goff, B. A., U. Hermanto, J. Rumbaugh, J. Blake, M. Bamberg and T. Hasan (1994) Photoimmunotherapy and biodistribution with an OC125-chlorin immunoconjugate in an *in vivo* murine ovarian cancer model. *Br. J. Cancer* **70**, 474-480.
- Goff, B. A., J. Blake, M. P. Bamberg and T. Hasan (1996) Treatment of ovarian cancer with photodynamic therapy and immunoconjugates in a murine ovarian cancer model. *Br. J. Cancer* **74**, 1194-1198.
- Goldman, M. J. and J. M. Wilson (1995) Expression of $\alpha_v\beta_5$ integrin is necessary for efficient adenovirus-mediated gene transfer in the human airway. *J. Virol.* **69**, 5951-5958.
- Goldsmith, S. J. (1997) Receptor imaging: competitive or complementary to antibody imaging? *Sem. Nucl. Med.* **27**, 85-93.
- Goldstein, J. L. and M. S. Brown (1974) Binding and degradation of low density lipoproteins by cultured human fibroblasts. *J. Bio. Chem.* **249**, 5153-5162.
- Goldstein, J. L., R. G. W. Anderson and M. S. Brown (1979) Coated pits, coated vesicles and receptor-mediated endocytosis. *Nature* **279**, 679-685.
- Goldstein, J. L., S. K. Basu and M. S. Brown (1983) Receptor-mediated endocytosis of low-density lipoprotein in cultured cells. *Meth. In Enzymol.* **98**, 241-260.
- Goligorsky, M. S., E. Noiri, H. Kessler and V. Romanov (1997) Therapeutic potential of RGD peptides in acute renal injury. *Kidney Int.* **51**, 1487-1492.

- Gómez, M., A. Luque, M. A. del Pozo, N. Hogg, F. Sánchez-Madrid and C. Cabañas (1997) Functional relevance during lymphocyte migration and cellular localization of activated $\beta 1$ integrins. *Eur. J. Immunol.* **27**, 8-16.
- Gorman, A. A. and M. A. J. Rodgers (1992) Current perspectives of singlet oxygen detection in biological environments. *J. Photochem. Photobiol. B: Biol.* **14**, 159-176.
- Granville, D. J., C. M. Carthy, D. W. Hunt and B. M. McManus (1998) Apoptosis: Molecular aspects of cell death and disease. *Lab. Invest.* **78**, 893-913.
- Granville, D. J., H. Jiang, M. T. An, J. G. Levy, B. M. McManus and D. W. C. Hunt (1998) Overexpression of Bcl-X_L prevents caspase-3-mediated activation of DNA fragmentation factor (DFF) produced by treatment with the photochemotherapeutic agent BPD-MA. *FEBS Lett.* **422**, 151-154.
- Granville, D. J., H. Jiang, M. T. An, J. G. Levy, B. M. McManus and D. W. Hunt (1999) Bcl-2 overexpression blocks caspase activation and downstream apoptotic events instigated by photodynamic therapy. *Br. J. Cancer* **79**, 95-100.
- Granville, D. J., J. R. Shaw, S. Leong, C. M. Carthy, P. Margaron, D. W. Hunt and B. M. McManus (1999) Release of cytochrome c, Bax migration, Bid cleavage and activation of caspases 2, 3, 6, 7, 8 and 9 during endothelial cell apoptosis. *Am. J. Pathol.* **155**, 1021-1025.
- Greber, U. F., M. Willets, P. Webster and A. Helenius (1993) Stepwise dismantling of adenovirus 2 during entry into cells. *Cell* **75**, 1-20.
- Greber, U. F., P. Webster, J. Weber and A. Helenius (1996) The role of the adenovirus protease in virus entry into cells. *EMBO* **15**, 1766-1777.
- Green, D. R. and J. C. Reed (1998) Mitochondria and apoptosis. *Science* **281**, 1309-1312.
- Grislain, L., P. Couvreur, V. Lenaerts, M. Roland, D. Deprez-Decampeneere and P. Speiser (1983) Pharmacokinetics and distribution of a biodegradable drug-carrier. *Int. J. Pharma.* **15**, 335-345.
- Gudgin Dickson, E. F., J. C. Kennedy and R. H. Pottier (2000) Protease-mediated delivery of light-activated drugs. *13th International Congress on Photobiology* July 1-6, 2000; San Francisco, California: Abstract 384.
- Gueddari, N., G. Favre, H. Hachem, E. Marek, F. Le Gaillard and G. Soula (1993) Evidence for up-regulated low density lipoprotein receptor in human lung adenocarcinoma cell line A549. *Biochimie* **75**, 811-819.
- Gueddari, N. G. Favre, C. Marmouget, G. Soula and F. Le Gaillard (1996) Involvement of tyrosine kinase activity in the low-density lipoprotein receptor expression in human lung adenocarcinoma cell line A549. *Biochimie* **78**, 874-881.
- Gupta, S., N. Ahmad and H. Mukhtar (1998) Involvement of nitric oxide during phthalocyanine (Pc4) photodynamic therapy-mediated apoptosis. *Cancer Res.* **58**, 1785-1788.
- Haas, T. A. and E. F. Plow (1994) Integrin-ligand interactions: a year in review. *Curr. Opinion Cell Biol.* **6**, 656-662.
- Hackbarth, S., V. Horneffer and B. Röder (2000) Multiple substituted dendrimers as multipliers in modular carrier-systems for PDT. *13th International Congress on Photobiology* July 1-6, 2000; San Francisco, California: Abstract 386.
- Haddada, H., L. Cordier and M. Perricaudet (1995) Gene therapy using adenovirus vectors. *Curr. Top. Microbiol. Immunol.* **199**, 97-306.
- Hadjur, C., G. Wagnières, F. Ihringer, Ph. Monnier and H van den Bergh (1997) Production of the free radical O₂⁻ and [•]OH by irradiation of the photosensitizer zinc(II) phthalocyanine. *J. Photochem. Photobiol. B: Biol.* **38**, 196-202.

- Hadjur, C., G. Wagnieres, P. Monnier and H. van den Bergh (1997) EPR and spectrophotometric studies of free radicals (O_2^- , $^{\bullet}OH$, BPD- $MA^{\bullet-}$) and singlet oxygen (1O_2) generated by irradiation of benzoporphyrin derivative monoacid ring A. *Photochem. Photobiol.* **65**, 818-827.
- Hahn, K. A., M. Panjehpour and X. Lu (1996) Fluence-dependent induction of micronuclei by aluminum phthalocyanine tetrasulfonate-mediated photodynamic therapy. *Photochem. Photobiol.* **63**, 117-122.
- Hahn, S. M. and E. Glatstein (1999) The emergence of photodynamic therapy as a major modality in cancer treatment. *Rev. Contemp. Pharmacother.* **10**, 69-74.
- Hamblin, M. R. and E. L. Newman (1994a) On the mechanism of tumour-localising effect in photodynamic therapy. *J. Photochem. Photobiol. B: Biol.* **23**, 3-8.
- Hamblin, M. R. and E. L. Newman (1994b) Photosensitizer targeting in photodynamic therapy I. Conjugates of haematoporphyrin with albumin and transferrin. *J. Photochem. Photobiol. B: Biol.* **26**, 45-56.
- Hamblin, M. R. and E. L. Newman (1994c) Photosensitizer targeting in photodynamic therapy II. Conjugates of haematoporphyrin with serum lipoproteins. *J. Photochem. Photobiol. B: Biol.* **26**, 147-157.
- Hamblin, M. R., J. L. Miller and T. Hasan (1996) Effect of charge on the interaction of site-specific photoimmunoconjugates with human ovarian cancer cells. *Cancer Res.* **56**, 5205-5210.
- Hamblin, M. R., M. Rajadhyaksha, T. Momma, N. S. Soukos and T. Hasan (1998) In vitro fluorescence imaging of the transport of charged chlorin e6 conjugates in a rat orthotopic prostate tumour. *Br. J. Cancer* **81**, 261-268.
- Hamblin, M. R., D. del Governatore, I. Rizvi and T. Hasan (2000) Biodistribution of charged 17.1A photoimmunoconjugates in a murine model of hepatic metastasis of colorectal cancer. *Br. J. Cancer* **83**, 1544-1551.
- Hamblin, M. R., J. L. Miller and B. Ortel (2000a) Scavenger-receptor targeted photodynamic therapy. *Photochem. Photobiol.* **72**, 533-540.
- Hamblin, M. R., M. del Governatore, C. R. Shea, I. Rizvi, K. L. Molpus and T. Hasan (2000) Experimental photoimmunotherapy for treatment of hepatic metastases of colorectal cancer in the mouse. *13th International Congress on Photobiology* July 1-6, 2000; San Francisco, California: Abstract 172.
- Hamblin, M. R., M. Huzaira and D. A. O'Donnell (2000) Scavenger receptor-targeted photodynamic therapy of murine tumors *in vivo*. *13th International Congress on Photobiology* July 1-6, 2000; San Francisco, California: Abstract 173.
- Hamblin, M. R., D. A. O'Donnell, N. Murthy, K. Rajagopalan, M. E. Sherwood and T. Hasan (2000) Photoinactivation of antibiotic-resistant bacteria using polycationic photosensitizer conjugates. *13th International Congress on Photobiology* July 1-6, 2000; San Francisco, California: Abstract 392.
- Hamblin, M. R., D. A. O'Donnell, N. Murthy, C. H. Contag, A. T. MacManus and T. Hasan (2000) Photodynamic inactivation of luminescent bacteria in contaminated and infected wounds in the mouse. *13th International Congress on Photobiology* July 1-6, 2000; San Francisco, California: Abstract 393.
- Hanahan, D. and J. Folkman (1996) Patterns and emerging mechanisms of the angiogenic switch during tumorigenesis. *Cell* **86**, 353-364.
- Harriman, A., G. Luengo and K. S. Gulliya (1990) In vitro photodynamic activity of kryptocyanine. *Photochem. Photobiol.* **52**, 735-740.
- Hasan, T., C. W. Lin and A. Lin (1989) Laser-induced selective cytotoxicity using monoclonal antibody-chromophore conjugates. *Prog. Clin. Biol. Res.* **288**, 471-477.
- Hasan, T. and J. A. Parrish. (1997) Photodynamic therapy of cancer. In *Cancer Medicine* (Eds. J. F. Holland, R. C. Bast, D. L. Morton, E. Frei III, D. W. Kufe and R. R. Weichselbaum) William and Wilkins, Baltimore; 4th ed, pp.739-751.
- Hausmann, W. (1908a) Die sensibilisierende wirkung tierischer farbstoffe und ihre physiologische bedeutung. *Wien. Klin Wchnschr.* **21**, 1527-1529.

- Hausmann, W. (1908b) Über die giftige wirkung der hämatoporphyrins auf warmblüter bei belichtung. *Wien. Klin. Wchnschr.* **22**, 1820-1821.
- Hausmann, W. (1911) Die sensibilisierende wirkung des hämatoporphyrins. *Biochem. Ztschr.* **30**, 276-316.
- Hautanen, A., J. Gailit, D. M. Mann and E. Ruoslahti (1989) Effects of modifications of the RGD sequence and its context on recognition by the fibronectin receptor. *J. Biol. Chem.* **264**, 1437-1442.
- He, X.-Y., R. A. Sikes, S. Thomsen, L. W. K. Chung and S. L. Jacques (1994) Photodynamic therapy with photofrinII induces programmed cell death in carcinoma cell lines. *Photochem. Photobiol.* **59**, 468-473.
- He, J., H. E. Larkin, Y.-S. Li, B. D. Rihter, S. I. A. Zaidi, M. A. J. Rodgers, H. Mukhtar, M. E. Kenney and N. L. Oleinick (1997) The synthesis, photophysical and photobiological properties and in vivo structure-activity relationships of a set of silicon phthalocyanine PDT photosensitizers. *Photochem. Photobiol.* **65**, 581-586.
- He., J., C. M. Whitacre, L.-Y. Xue, N. A. Berger and N. L. Oleinick (1998) Protease activation and cleavage of poly(ADP-ribose) polymerase: An integral part of apoptosis in response to photodynamic treatment. *Cancer Res.* **58**, 940-946.
- He., J., M.-F. Horng, J. T. Deahl, N. L. Oleinick and H. H. Evans (1998) Variations in photodynamic efficacy during the cell, 720-728. ular uptake of two phthalocyanine photosensitizers. *Photochem. Photobiol.* **67**, 720-728.
- Hecht, S. S., S. Isaacs and N. Trushin (1994) Lung tumor induction in A/J mice by the tobacco smoke carcinogens 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone and benzo[a]pyrene: a potentially useful model for evaluation of chemopreventive agents. *Carcinogenesis* **15**, 2721-2725.
- Henderson, B. W. and A. C. Miller (1986) Effects of scavengers of reactive oxygen and radical species on cell survival following photodynamic treatment *in vitro*: Comparison to ionizing radiation. *Radiat. Res.* **108**, 196-205.
- Henderson, B. W. and G. Farrell (1989) Possible implications of vascular damage for tumor cell inactivation in vivo: Comparison of different photosensitizers. *SPIE Conf. Proc.* **1065**, 2-10.
- Henderson, B. W. and E. Mayhew (1990) Experience with the liposomal delivery of the photosensitizer isoBoSiNc. *Proc. Int. Soc. Optical Eng.* **1203**, 126-135.
- Herisse, J. and F. Galibert (1981) Nucleotide sequence of the EcoRI E fragment of adenovirus 2 genome. *Nucl. Acids Res.* **9**, 1229-1240.
- Hirasawa, M., N. Shijubo, T. Uede and S. Abe (1994) Integrin expression and ability to adhere to extracellular matrix proteins and endothelial cells in human lung cancer lines. *Br. J. Cancer* **70**, 466-473.
- Ho, Y. K., R. G. Smith, M. S. Brown and J. L. Goldstein (1978) Low density lipoprotein (LDL) receptor activity in human acute myelogenous leukemia cells. *Blood* **52**, 1099-1104.
- Hoebcke, M. (1995) The importance of liposome as models and tools in the understanding of photosensitization mechanisms. *J. Photochem. Photobiol. B: Biol.* **28**, 189-196.
- Hollowood, A. D., T. Lai, C. M. Perks, P. V. Newcomb, D. Alderson and J. M. P. Holly (2000) IGFBP-3 prolongs the p53 response and enhances apoptosis following UV irradiation. *Int. J. Cancer* **88**, 336-341.
- Hong, S. S. and P. Boulanger (1995) Protein ligands of the human adenovirus type 2 outer capsid identified by biopanning of a phage-displayed peptide library on separate domains of wild-type and mutant penton capsomers. *EMBO* **14**, 4714-4727.
- Hopkins, CR. (1986) Site-specife drug delivery-cellular opportunities and changes. In *Site-specific drug delivery* (Eds. E. Tomlinson and S. S. Davis) John Wiley & Sons Ltd, New York, pp.27-48.
- Hosono, M. N., M. Hosono, K. Endo, R. Ueda and Y. Onoyama (1994) Effect of hyperthermia on tumor uptake of radiolabeled anti-neural cell adhesion molecule antibody in small-cell lung cancer xenografts. *J. Nuclear Med.* **35**, 504-509.

- Hsi, R. A., D. I. Rosenthal and E. Glatstein (1999) Photodynamic therapy in the treatment of cancer. *Drugs* **57**, 725-734.
- Huang, S., R. I. Endo and G. R. Nemerow (1995) Upregulation of integrins $\alpha_v\beta_3$ and $\alpha_v\beta_5$ on human monocytes and T lymphocytes facilitates adenovirus-mediated gene delivery. *J. Virol.* **69**, 2257-2263.
- Huang, S., T. Kamata, Y. Takada, Z. M. Ruggeri and G. R. Nemerow (1996) Adenovirus interaction with distinct integrins mediates separate events in cell entry and gene delivery to hematopoietic cells. *J. Virol.* **70**, 4502-4508.
- Huttenlocher, A., R. R. Sandborg and A. F. Horwitz (1995) Adhesion in cell migration. *Curr. Opinion Cell Biol.* **7**, 697-706.
- Isele, U., K. Schieweck, R. Kessler, P. van Hoogevest and H.-G. Capraro (1995) Pharmacokinetics and body distribution of liposomal zinc phthalocyanine in tumor-bearing mice: Influence of aggregation state, particle size, and composition. *J. Pharmaceut. Sci.* **84**, 166-173.
- Iwamoto, H., H. Sakai, S. Tada, M. Nakamuta and H. Nawata (1999) Induction of apoptosis in rat hepatic stellate cells by disruption of integrin-mediated cell adhesion. *J. Lab. Clin. Med.* **134**, 83-89.
- Jasanada, F., P. Urizzi, J.-P. Souchard, F. Le Gaillard, G. Favre and F. Nepveu (1996) Indium-111 labeled low density lipoprotein (LDL) with the DTPA-bis(stearylamide): evaluation as a potential radiopharmaceutical for tumor localization. *Bioconj. Chem.* **7**, 72-81.
- Jiang, F. N., S. Jiang, D. Liu, A. Richter, J. G. Levy (1990) Development of technology for linking photosensitizers to a model monoclonal antibody. *J. Immunol. Methods* **134**, 139-149.
- Jiang, F. N., D. J. Liu, H. Neyndorff, M. Chester, S.-Y. Jiang and J. G. Levy (1991) Photodynamic killing of squamous cell carcinoma cells using a monoclonal antibody-photosensitizer conjugate. *J. Natl. Cancer Inst.* **83**, 1218-1225.
- Jiao, D., K. E. Eklind, C.-I. Choi, D. H. Desai, S. G. Amin and F.-L. Chung (1994) Structure-activity relationships of isothiocyanates as mechanism-based inhibitors of 4-(methylnitrosamino)-1-(2-pyridyl)-1-butanone-induced lung tumorigenesis in A/J mice. *Cancer Res.* **54**, 4327-4333.
- Johnson, R., L. Staiano-Coico, L. Austin, I. Cardinale, R. Nabeya-Tsukifuji and J. G. Krueger (1996) PUVA treatment selectively induces a cell cycle block and subsequent apoptosis in human T-lymphocytes. *Photochem. Photobiol.* **63**, 566-571.
- Jones, E. Y. (1996) Three-dimensional structure of cell adhesion molecules. *Curr. Opinion Cell Biol.* **8**, 602-608.
- Jori, G., M. Beltramini, E. Reddi, B. Salvato, A. Pagnan, L. Ziron, L. Tomio and T. Tsanov (1984) Evidence for a major role of plasma lipoprotein as hematoporphyrin carriers *in vivo*. *Cancer Lett.* **24**, 291-297.
- Jori, G. and E. Reddi (1993) The role of lipoproteins in the delivery of tumour-targeting photosensitizers. *Int. J. Biochem.* **25**, 1369-1375.
- Jori, G. and C. Fabris (1998) Relative contributions of apoptosis and random necrosis in tumour response to photodynamic therapy: effect of the chemical structure of Zn(II)-phthalocyanine. *J. Photochem. Photobiol. B: Biol.* **43**, 181-185.
- Jornvall, H., G. Akusjarvi, P. Alestrom, H. von Bahr-Lindstrom, U. Pettersson, E. Appella, A. V. Fowler and L. Philipson (1981) The adenovirus hexon protein. The primary structure of the polypeptide and its correlation with the hexon gene. *J. Biol. Chem.* **256**, 6181-6186.
- Juliano, R. L. and J. A. Varner (1993) Adhesion molecules in cancer: the role of integrins. *Curr. Opinion Cell Biol.* **5**, 812-818.
- Kanofsky, J. R. (1991) Quenching of singlet oxygen by human red cell ghosts. *Photochem. Photobiol.* **53**, 93-99.

Gay, J. Gerfaux and P. A. Boulanger (1994) Oligomerization of recombinant penton base 2 and its assembly with fiber in baculovirus-infected cells. *Virology* **202**, 782-795.

Letokhov (1994) Possible benefits of two-quantum excitation in ALA-PDT? *J. Photochem. Biol.* **23**, 261-262.

i) Porphyrin-lipoprotein association as a factor in porphyrin localization. *Cancer Lett.* **33**,

ompson, K. Saatio and K. D. Natwi. (1987) Tumor localization and photosensitization by derivatives of tetraphenylporphine. *Photochem. Photobiol.* **45**, 787-790.

forjan and G. M. Garbo (1991) Sites and efficacy of photodamage by tin etiopurpurin in parent delivery systems. *Photochem. Photobiol.* **54**, 193-196.

ii) Interactions between N-aspartyl chlorin e6, detergent micelles and plasma lipoproteins. *Photobiol.* **61**, 646-649.

Y. Luo (1996) Delayed oxidative photodamage induced by photodynamic therapy. *Photobiol.* **64**, 601-604.

Luo, Y. Deng and C. K. Chang (1997) The role of subcellular localization in initiation of photodynamic therapy. *Photochem. Photobiol.* **65**, 422-426.

iii) T. J. Dougherty (1999) Agents used in photodynamic therapy. *Rev. Contemp. Phys.* **10**, 19-24.

Y. Luo (1999) Photodynamic therapy: A mitochondrial inducer of apoptosis. *Cell Death and Dis.* **10**, 28-35.

Y. Luo (2001) Intracellular sites of photodamage as a factor in apoptotic cell death. *J. Photochem. Photobiol.* **5**, 181-184.

MacRoberts, P. M. Speight and J. H. Bennett (1998) Induction of apoptotic cell death by photodynamic therapy in human keratinocytes. *Arch. Oral Biol.* **43**, 143-149.

A. Veloso Jr, F. Tolentino, T. Hasan and M. R. Hamblin (1999) Photodynamic tissue damage by chlorin e6 protein conjugates. *Invest. Ophthalmol. Vis. Sci.* **40**, 3132-3137.

iv) C. Breistol and Ø. Fodstad (1992) Site-dependent differences in sensitivity of LOX human tumors in nude rats to dacarbazine and mitoxantrone, but not to doxorubicin and cisplatin. *Int. J. Cancer* **51**, 1347-1451.

v) B. A., I. S. Nechaeva and G. V. Ponomarev (1994) Approaches to targeted photodynamic therapy. *J. Controlled Release* **29**, 1-16.

vi) Wang and E. Ruoslahti (1994) Isolation of a highly specific ligand for the $\alpha_5\beta_1$ integrin using a phage display library. *J. Cell Biol.* **124**, 373-380.

vii) B. Seed (1997) Integrins and inside-out signal transduction: converging signals from PKC and Ras. *Opinion in Cell Biol.* **9**, 725-731.

viii) J. Moan and S. B. Brown (1989) The distribution of porphyrins with different tumor localization among human plasma proteins. *Br. J. Cancer* **59**, 184-188.

ix) S. Zhao, Z. Xie, H. Segall, A. Younes, D. F. Claxton, Z. Estrov, S. M. Kornblau and M. J. Scher (1993) Apoptosis, molecules and mechanisms. *Drug Resistance in Leukemia and Lymphoma III*. Kluwer Academic/Plenum Publishers, New York, pp. 217-236.

x) N. L. Krinick, B. Rihova and K. Ulbrich (1990) Targetable N-(2-methacrylamide copolymer-chlorin e6 conjugates. *SPIE Conf. Proc.* **1203**, 144-152.

xi) (1993) Cellular delivery and retention of Photofrin: III. Role of plasma proteins in clearance from cells. *Photochem. Photobiol.* **57**, 846-850.

- Korbelik, M. and G. Krosi (1994) Enhanced macrophage cytotoxicity against tumor cells treated with photodynamic therapy. *Photochem. Photobiol.* **60**, 497-502.
- Korbelik, M. and G. Krosi (1995) Photofrin accumulation in malignant and host cell populations of a murine fibrosarcoma. *Photochem. Photobiol.* **62**, 162-168.
- Krasnovsky, A. A. (1981) Quantum yield of photosensitized luminescence and radiative lifetime of singlet ($^1\Sigma_g^+$) molecular oxygen in solutions. *Chem. Phys. Lett.* **81**, 443-445.
- Krasnykh, V. N., G. V. Mikheeva, J. T. Douglas and D. T. Curiel (1996) Generation of recombinant adenovirus vectors with modified fibers for altering viral tropism. *J. Virol.* **70**, 6839-6846.
- Krieg, M., H. Weisser and S. Tunn (1995) Potent activities of androgen metabolizing enzymes in human prostate. *J. Steroid Biochem. Mol. Biol.* **53**, 395-400.
- Krieger, M., S. Acton, J. Ashekenas, A. Pearson, M. Penman and D. Resnick (1993) Molecular flypaper, host defense and atherosclerosis. Structure, binding properties and functions of macrophage scavenger receptors. *J. Biol. Chem.* **268**, 4569-4572.
- Krinick, N. L., Y. Sun, D. A. Joyner, R. Reed, J. D. Spikes, R. C. Straight and J. Kopecek (1992) Polymer-bound meso-chlorin e6 for PDT. *SPIE Conf. Proc.* **1645**, 142-154.
- Krinick, N. L., Y. Sun, D. A. Joyner, J. D. Spikes, R. C. Straight, and J. Kopecek (1994) A polymeric drug delivery system for the simultaneous delivery of drugs activatable by enzymes and/or light. *J. Biomater. Sci. Polym. Ed.* **5**, 303-324.
- Labib, A., V. Lenaerts, F. Chouinard, J. C. Leroux, R. Ouellet and J. E. van Lier (1991) Biodegradable nanospheres containing phthalocyanines and naphthalocyanines for targeted photodynamic tumor therapy. *Pharma. Res.* **8**, 1027-1031.
- Lam, S. (1994) Photodynamic therapy of lung cancer. *Sem. Oncol.* **21**, 15-19.
- Langlois, R., H. Ali, N. Brasseur, J. R. Wagner and J. E. van Lier (1986) Biological activities of phthalocyanines IV. Type II sensitized photooxidation of L-tryptophan and cholesterol by sulfonated metallophthalocyanines. *Photochem. Photobiol.* **44**, 117-123.
- Langlois, R., H. Ali and J. E. van Lier (1993) Réaction du L-tryptophan avec le radical peroxyde du trichlorocarbène ($\text{CCl}_3\text{OO}\cdot$) et l'oxygène singulet ($^1\text{O}_2$): Formation de différentes paires d'hydroperoxydes isomériques. *J. Chim. Phys.* **90**, 985-999.
- Lapointe, D., N. Brasseur, J. Cadorette, C. La Madeleine, S. Rodrique, J. E. van Lier and R. Lecomte (1999) High-resolution PET imaging for in vivo monitoring of tumor response after photodynamic therapy in mice. *J. Nucl. Med.* **40**, 876-882.
- Larroque, C., A. Pelegrin and J. E. van Lier (1996) Serum albumin as a vehicle for zinc phthalocyanine: Photodynamic activities in solid tumour models. *Br. J. Cancer* **74**: 1886-1890.
- Latina, M. A., P. H. Kobsa, S. L. Rakestraw, E. A. Cren, T. Hasan and M. L. Yarmush (1988) Photochemical targeting of phagocytic trabecular meshwork cells using chlorin e6 coupled microsphere. *Proc. Soc. Photo-Optic Instr. Eng.* **997**, 32-41.
- Laukka, M. A., K. K. Wang and J. A. Bonner (1994) Apoptosis occurs in lymphoma cells but not in hepatoma cells following ionizing radiation and photodynamic therapy. *Digestive Diseases and Sci.* **39**, 2467-2475.
- Lawrence, D.S. and D. G. Whitten (1996) Photochemistry and photophysical properties of novel, unsymmetrically substituted metallophthalocyanines. *Photochem. Photobiol.* **64**, 923-935.
- Lee, S. K., I. J. Forbes and W. H. Betts (1984) Oxygen dependency of photocytotoxicity with hematoporphyrin derivative. *Photochem. Photobiol.* **39**, 631-634.
- Lembert, N. and L.-Å. Idahl (1995) Regulatory effects of ATP and luciferin on firefly luciferase activity. *Biochem. J.* **305**, 929-933.

- Lenaerts, V., A. Labib, F. Chouinard, J. Rousseau, H. Ali and J. E. van Lier (1995) Nanocapsules with a reduced liver uptake: Targeting of phthalocyanines to EMT-6 mouse mammary tumour in vivo. *Eur. J. Pharm. Biopharm.* **41**, 38-43.
- Lestavel-Delattre S., F. Martin-Nizard, V. Clavey, P. Testard, G. Favre, G. Dounalin, H. S. Houssaini, J.-M. Bard, P. Duriez, C. Delbart, G. Soula, D. Lesieur, I. Lesieur, J.-C. Cazin, M. Cazin and J.-C. Fruchard (1992) Low-density lipoprotein for delivery of an acrylophenone antineoplastic molecule into malignant cells. *Cancer Res.* **52**, 3629-3635.
- Levy, J. G. (1994) Photosensitizers in photodynamic therapy. *Sem. Oncol.* **21**, 4-10.
- Levy, J. G. (1995) Photodynamic therapy. *Tibtech.* **13**, 14-18.
- Leznoff, C. C. and A. B. P. Lever (1996) Series Preface. in *Phthalocyanines: Properties and Applications Volume 4* (eds. C. C. Leznoff and A. B. P. Lever). New York: VCH Publishers, Inc., pp. v.
- Li, Y., B. Lin, A. Agadir, R. Liu, M. I. Dawson, J. C. Reed, J. A. Fontana, F. Bost, P. D. Hobbs, Y. Zheng, G. Chen, B. Shroot, D. Mercola and X. Zhang (1998) Molecular determinants of AHPN (CD437)-induced growth arrest and apoptosis in human lung cancer cell lines. *Molecular and Cellular Biol.* **18**, 4719-4731.
- Linstead, R. P. (1934) Phthalocyanines. Part I. A new type of synthetic colouring matters. *J. Chem. Soc.* 1016-1017.
- Linstead, R. P. and A. R. Lowe (1934a) Phthalocyanines. Part III. Preliminary experiments on the preparation of phthalocyanines from phthalonitrile. *J. Chem. Soc.* 1022-1027.
- Linstead, R. P. and A. R. Lowe (1934b) Phthalocyanines. Part V. The molecular weight of magnesium phthalocyanine. *J. Chem. Soc.* 1031-1033.
- Linstead, R. P. and J. M. Robertson (1936) The stereochemistry of metallic phthalocyanines. *J. Chem. Soc.* 1736-1738.
- Lipson, R. L., E. J. Baldes and E. M. Olsen (1961) The use of a derivative of hematoporphyrin in tumor detection. *J. Natl. Cancer Inst.* **26**, 1-11.
- Lister-James, J., L. C. Knight, A. H. Maurer, L. R. Bush, B. R. Moyer and R. T. Dean (1995) Thrombus imaging with a technetium-99m-labeled activated platelet receptor-binding peptide. *J. Nucl. Med.* **37**, 775-781.
- Lister-James, J., B. R. Moyer and R.T. Dean (1996) Small peptides radiolabeled with ^{99m}Tc. *Q.J. Nucl. Med.* **40**, 221-233.
- Lister-James, J., B. R. Moyer and R.T. Dean (1997) Pharmacokinetic considerations in the development of peptide-based imaging agents. *Q. J. Nucl. Med.* **41**, 111-118.
- Lister-James, J., S. Vallabhajosula, B. R. Moyer, D. A. Pearson, B. J. McBride, M. A. De Rosch, L. R. Bush, J. Machac and R. T. Dean (1997) Pre-clinical evaluation of a technetium-99m platelet receptor-binding peptide. *J. Nucl. Med.* **38**, 105-111.
- Lombardi, P., G. Norata, F. M. Maggi, G. Canti, P. Franco, A. Nicolin and A. L. Catapano (1989) Assimilation of LDL by experimental tumors in mice. *Photochem. Photobiol.* **1003**, 301-306.
- Louis, N., P. Fender, A. Barge, P. Kitts and J. Chroboczek (1994) Cell-binding domain of adenovirus serotype 2 fiber. *J. Virol.* **68**, 4104-4106.
- Love, W. G., S. Duk, R. Biolo, G. Jori and P. W. Taylor (1996) Liposome-mediated delivery of photosensitizers: Localization of zinc (II)-phthalocyanine within implanted tumors after intravenous administration. *Photochem. Photobiol.* **63**, 656-661.
- Lu, J. M., C. M. Peterson, J. G. Shiah, Z. W. Gu, C. A. Peterson, R. C. Straight and J. Kopecek (1999) Cooperativity between free and N-(2-hydroxypropyl) methacrylamide copolymer bound adriamycin and meso-chlorin e6 monoethylenediamine induced photodynamic therapy in human epithelial ovarian carcinoma in vitro. *Int. J. Oncol.* **15**, 5-16.

- Lukyanets, E. A. (1999) Phthalocyanines as photosensitizers in the photodynamic therapy of cancer. *J. Porphyrins Phthalocyanines* **3**, 424-432.
- Luo, Y. and D. Kessel (1996) The phosphatase inhibitor calyculin antagonizes the rapid initiation of apoptosis by photodynamic therapy. *Biochem. Biophys. Res. Comm.* **221**, 72-76.
- Luo, Y., C. K. Chang and D. Kessel (1996) Rapid initiation of apoptosis by photodynamic therapy. *Photochem. Photobiol.* **63**, 528-534.
- Luo, Y. and D. Kessel (1997) Initiation of apoptosis versus necrosis by photodynamic therapy with chloroaluminum phthalocyanine. *Photochem. Photobiol.* **66**, 479-483.
- Lutsenko, S. V., N. B. Feldman, G. V. Finakova, G. A. Posypanova, S. E. Severin, K. G. Skryabin, M. P. Kirpichnikov, E. A. Lukyanets and G. N. Vorozhtsov (1999) Targeting phthalocyanines to tumor cells using epidermal growth factor conjugates. *Tumor Biol.* **20**, 218-224.
- Macdonald, I. J. and T. J. Doherty (2001) Basic principles of photodynamic therapy. *J. Porphyrins and Phthalocyanines* **5**, 105-129.
- MacRoberts, A. J., S. G. Brown and D. Phillips (1989) What are the ideal photoproperties for a sensitizer? In. *Photosensitizing Compounds: Their Chemistry, Biology and Clinical Use*. Ciba Foundation Symposium 146. UK: John Wiley and Sons, pp. 4-12.
- Maman, N., S. Dhami, D. Phillips and D. Brault (1999) Kinetic and equilibrium studies of incorporation of di-sulfonated aluminum phthalocyanine into unilamellar vesicles. *Biochim. Biophys. Acta* **1420**, 168-178.
- Marcon, N. E. (1994) Photodynamic therapy and cancer of the esophagus. *Sem. Oncol.* **21**, 20-23.
- Margalit, R. and E. Silbiger (1985) Albumin microspheres as delivery systems for photodynamic drugs: physico-chemical studies and their implications for in vivo situations. *J. Microencapsulation* **2**, 183-196.
- Margaron, P., R. Langlois, J. E. van Lier and S. Gaspard (1992) Photodynamic properties of naphthalosulfobenzoporphyrines, novel asymmetric, amphiphilic phthalocyanine derivatives. *J. Photochem. Photobiol. B: Biol.* **14**, 187-199.
- Margaron, P. P. Madarnas, R. Ouellet and J. E. van Lier (1996) Biological activities of phthalocyanines. XVII Histopathologic evidence for different mechanisms of EMT-6 tumor necrosis induced by photodynamic therapy with disulfonated aluminum phthalocyanine or Photofrin*. *Anticancer Res.* **16**, 613-620.
- Margaron, P., M. Gregoire, V. Scasnar, H. Ali and J. E. van Lier (1996) Structure-photodynamic activity relationships for a series of 4-substituted phthalocyanines. *Photochem. Photobiol.* **63**, 217-223.
- Martsev, S. P., V. A. Preygerzon, Y. I. Mel'nikova, Z. I. Kravchuk, G. V. Ponomarev, V. E. Lunev and A. P. Savitsky (1995) Modification of monoclonal and polyclonal IgG with palladium(II) coproporphyrin I: Stimulatory and inhibitory functional effects induced by two different methods. *J. Immunol. Methods* **186**, 293-304.
- Mathias, P., T. Wickham, M. Moore and G. Nemerow (1994) Multiple adenovirus serotypes use α_v integrins for infection. *J. Virol.* **68**, 6811-6814.
- Matthews, W., J. Cook, J. B. Mitchell, R. R. Perry, S. Evans and H. I. Pass (1989) *In vitro* photodynamic therapy of lung cancer: investigation of dose-rate effects. *Cancer Res.* **49**, 1718-1721.
- Mattos, C. and D. Ringe (1996) Locating and characterizing binding sites on proteins. *Nature Biotech.* **14**, 595-599.
- Mazière, J. C., P. Molière and R. Santus (1991) The role of the low density lipoprotein receptor pathway in the delivery of lipophilic photosensitizers in the photodynamic therapy of tumours. *J. Photochem. Photobiol. B: Biol.* **8**, 351-360.
- McCaughan, J. S. (1999) Photodynamic therapy. A review. *Drugs and Aging* **15**, 49-68.

- McCray, P. B., K. Armstrong, K., J. Zabner, D. W. Miller, G. A. Koretzky, L. Couture, J. E. Robillard, A. E. Smith and M. J. Welsh (1995) Adenoviral-mediated gene transfer to fetal pulmonary epithelia in vitro and in vivo. *J. Clin. Invest.* **95**, 2620-2632.
- McCubbin, I. and D. Phillips (1986) The photophysics and photostability of zinc(II) and aluminium(III) sulphonated naphthalocyanines. *J. Photochem.* **34**, 187-195.
- Ménard, I. (1998) Vectorization des phthalocyanines par un anticorps monoclonal. M. Sc. Thesis, Université de Sherbrooke, Sherbrooke, Québec, Canada.
- Meredith, J. E. Jr. and M. A. Schwartz (1997) Integrins, adhesion and apoptosis. *Trends in Cell Biol.* **7**, PAGE NUMBER!
- Mew, D., C.-K. Wat, G. H. N. Towers and J. G. Levy (1983) Photoimmunotherapy: Treatment of animal tumors with tumor-specific monoclonal antibody-hematoporphyrin conjugates. *J. Immunol.* **130**, 1473-1477.
- Mew, D., V. Lum, C.-K. Wat, G. H. N. Towers, C.-H. C. Sun, R. J. Walter, W. Wright, M. W. Berns and J. G. Levy (1985) Ability of specific monoclonal antibodies and conventional antisera conjugated to hematoporphyrin to label and kill selected cell lines subsequent to light activation. *Cancer Res.* **45**, 4380-4386.
- Meyer-Betz, F. (1913) Untersuchungen über die biologische (photodynamische) wirkung des hämatoporphyrins und anderer derivative des blute und gallenfarbstoffs. *Deutsches Arch. Klin. Med.* **112**, 476-503.
- Miller, J. W., A. W. Walsh, M. Kramer, T. Hasan, N. Michaud, T. J. Flotte, R. Haimovici and E. Gragoudas (1995) Photodynamic therapy of experimental choroidal neovascularization using lipoprotein-delivered benzoporphyrin. *Arch. Ophthalmol.* **113**, 810-818.
- Miller, S. A., D. D. Dykes and H. F. Polesky (1988) A simple salting out procedure for extracting DNA from human nucleated cells. *Nuc. Acids Res.* **16**, 1215.
- Miquel, K., A. Pradines and G. Favre (1996) Farnesol and geranylgeraniol induce actin cytoskeleton disorganization and apoptosis in A549 lung adenocarcinoma cells. *Biochem. Biophys. Res. Comm.* **225**, 869-876.
- Miquel, K., A. Pradines, J. Sun, Y. Qian, A. D. Hamilton, S. M. Seti and G. Favre (1997) GGTI-298 induces G0-G1 block and apoptosis whereas FTI-277 causes G2-M enrichment in A549 cells. *Cancer Res.* **57**, 1846-1850.
- Misicka, A. and V. J. Hruby (1994) Optimization of disulfide bond formation. *Polish J. Chem.* **68**, 893-899.
- Mitchell, J. B., S. McPherson and W. DeGraff (1985) Oxygen dependence of hematoporphyrin derivative induced photoinactivation of Chinese hamster cells. *Cancer Res.* **45**, 2008-2011.
- Miyauchi, S., T. Imaoka, T. Utsunomiya, K. Hayashi, M. Kubo, T. Kawaguchi and Y. Matsui (1994) Oral administration of BOF-A2 to rats with lung transplanted tumors results in increased 5-fluorouracil levels. *Jpn. J. Cancer Res.* **85**, 665-668.
- Moan, J. and E. Boye (1981) Photodynamic effect on DNA and cell survival of human cells sensitized with hematoporphyrin. *Photobiochem. Photobiophys.* **2**, 301-307.
- Moan, J. (1990) On the diffusion length of singlet oxygen in cells and tissues. *J. Photochem. Photobiol. B: Biol.* **6**, 343-344.
- Montgomery, A. M. P., R. A. Reisfeld, and D. A. Cheresh (1994) Integrin $\alpha_v\beta_3$ rescues melanoma cells from apoptosis in three-dimensional dermal collagen. *Proc. Natl. Acad. Sci. USA.* **91**, 8856-8860.
- Moor, A. C., A. E. Wagenaar-van Gompel, A. Brand, T. M. A. R. Dubbelman and J. van Steveninck (1997) Primary targets for photoinactivation of vesicular stomatitis virus by AlPcS₄ or Pc4 and red light. *Photochem. Photobiol.* **65**, 465-470.

- Moor, A. C. E. (2000) Signaling pathways in cell death and survival after photodynamic therapy. *J. Photochem. Photobiol. B: Biol.* **57**, 1-13.
- Moor, A. C. E., I. Rizvi, M. Savellano, W. Ya, M. G. Del Carmen and T. Hasan (2000) Photoimmunotargeting of the epidermal growth factor receptor for the treatment of ovarian cancer. *13th International Congress on Photobiology* July 1-6, 2000; San Francisco, California: Abstract 507.
- Moore, J. V., C. M. L. West and C. Whitehurst (1997) The biology of photodynamic therapy. *Phys. Med. Biol.* **42**, 913-935.
- Moreno-Manzano, V., J. Lucio-Cazana, T. Konta, K. Nakayama and M. Kitamura (2000) Enhancement of TNF- α -induced apoptosis by immobilized arginine-glycine-aspartate: involvement of a tyrosine kinase-dependent, MAP kinase-independent mechanism. *Biochem. Biophys. Res. Comm.* **277**, 293-298.
- Morgan, A., G. M. Garbo, R. W. Keck and S. H. Selman (1988) New photosensitizers for photodynamic therapy: Combined effect of metalloporphyrin derivatives and light on transplantable bladder tumors. *Cancer Res.* **48**, 194-198.
- Morgan, J., A. G. Gray and E. R. Huehns (1989) Specific targeting and toxicity of sulphonated aluminium phthalocyanine photosensitised liposomes directed to cells by monoclonal antibody in vitro. *Br. J. Cancer* **59**, 366-370.
- Morgan, J., A. J. MacRobert, A. G. Gray and E. R. Huehns (1992) Use of photosensitive antibody directed liposomes to destroy target populations of cells in bone marrow: a potential purging method for autologous bone marrow transplantation. *Br. J. Cancer* **65**, 58-64.
- Morgan, J., H. Lottman, C. C. Abbou and D. K. Chopin (1994) A comparison of direct and liposomal antibody conjugates of sulfonated aluminum phthalocyanines for selective photoimmunotherapy of human bladder carcinoma. *Photochem. Photobiol.* **60**, 486-496.
- Moser, F. H and A. L. Thomas (1983) *The Phthalocyanines. Volume 1 and 2*. Boca Raton: CRC Press, Ltd.
- Mosqueira, V. C., P. Legrand, R. Gref, B. Heurtault, M. Appel and G. Barratt (1999) Interaction between macrophage cell line (J774A1) and surface-modified poly(D,L-lactide) nanocapsules bearing poly(ethylene glycol). *J. Drug Targeting* **7**, 65-78.
- Mosquera, V. C., P. Legrand, H. Pinto-Alphandary, F. Puisieux and G. Barratt (2000) Poly(D,L-lactide)nanocapsules prepared by a solvent displacement process: influence of the composition on physicochemical and structural properties. *J. Pharmaco. Sci.* **89**, 614-626.
- Mukhopadhyay, A., G. Chaudhuri, S. K. Arora, S. Sehgal and S. K. Basu (1989) Receptor-mediated drug delivery to macrophages in chemotherapy of Leishmaniasis. *Science* **244**, 705-707.
- Mukhopadhyay, T. and J. A. Roth (1997) Induction of apoptosis in human lung cancer cells after wild-type p53 activation by methoxestradiol. *Oncogene* **14**, 379-384.
- Murphy, S. T., J. Racz, C. S. Foote, C. M. Allen and J. E. van Lier (2001) Structure-photodynamic activity relationship of a series of amphiphilic silicon phthalocyanines with charged axial ligands. (in press)
- Muto, P., S. Lastoria, P. Varrella, E. Vergara, M. Salvatore, G. Morgano, J. Lister-James, J. D. Bernardy, R. T. Dean, D. Wencker and J. S. Borer (1995) Detecting deep venous thrombosis with technetium-99m-labeled synthetic peptide p280. *J. Nucl. Med.* **36**, 1384-1391.
- Nagae, T., A. Y. Louie, K. Aizawa, Sh. Ishimaru and S. E. Wilson (1998) Selective targeting and photodynamic destruction of intimal hyperplasia by scavenger-receptor mediated protein-chlorin e_6 conjugates. *J. Cardiovasc. Surg.* **39**, 709-715.
- Nicholson, D. W. (1999) Caspase structure, proteolytic substrates, and function during apoptotic cell death. *Cell Death and Differentiation* **6**, 1028-1042.
- Noodt, B. B., K. Berg, T. Stokke, Q. Peng and J. M. Nesland (1996) Apoptosis and necrosis induced with light and 5-aminolaevulinic acid-derived protoporphyrin IX. *Br. J. Cancer* **74**, 22-29.

- Novelli, A. and P. A. Boulanger (1991) Deletion analysis of functional domains in baculovirus-expressed adenovirus type 2 fiber. *Virology* **185**, 365-376.
- O'Reilly, M. S., L. Holmgren, Y. Shing, C. Chen, R. A. Rosenthal, M. Moses, W. S. Lane, Y. Cao, E. H. Sage and J. Folkman (1994) Angiostatin: A novel angiogenesis inhibitor that mediates the suppression of metastases by a lung lewis carcinoma. *Cell* **79**, 315-328.
- O'Reilly, M. S., L. Holmgren, C. Chen and J. Folkman (1996) Angiostatin induces and sustains dormancy of human primary tumors in mice. *Nature Med.* **2**, 689-692.
- O'Reilly, M. S., T. Boehm, Y. Shing, N. Fukai, G. Vasios, W. S. Lane, E. Flynn, J. R. Birkhead, B. R. Olsen and J. Folkman (1997) Endostatin: An endogenous inhibitor of angiogenesis and tumor growth. *Cell* **88**, 277-285.
- Obochi, M. O. K., R. W. Boyle and J. E. van Lier (1993) Biological activities of phthalocyanines. XIII. The effects of human serum components on the *in vitro* uptake and photodynamic activity of zinc phthalocyanine. *Photochem. Photobiol.* **57**, 634-640.
- Ochsner, M. (1996) Light scattering of human skin: a comparison between zinc (II)-phthalocyanine and photofrin II. *J. Photochem. Photobiol. B: Biol.* **32**, 3-9.
- Ochsner, M. (1997) Photophysical and photobiological processes in the photodynamic therapy of tumours. *J. Photochem. Photobiol. B: Biol.* **39**, 1-18.
- Oku, N., Y. Tokudome, C. Koike, N. Nishikawa, H. Mori, I. Saiki and S. Okada (1996) Liposomal ARG-GLY-ASP analogs effectively inhibit metastatic B16 melanoma colonization in murine lungs. *Life Sciences* **58**, 2263-2270.
- Oku, N., N. Saito, Y. Namba, H. Tsukada, D. Dolphin and S. Okada (1997) Application of long-circulating liposomes to cancer photodynamic therapy. *Biol. Pharm. Bull.* **20**, 670-673.
- Oku, N. (1999) Anticancer therapy using glucuronate modified long-circulating liposomes. *Adv. Drug Del. Rev.* **40**, 63-73.
- Okunaka, T. and H. Kato (1999) Potential applications of photodynamic therapy. *Rev. Contemp. Pharmacother.* **10**, 59-68.
- Oleinick, N. L., A. R. Antunez, M. E. Clay, B. D. Richter and M. E. Kenney (1993) New phthalocyanine photosensitizers for photodynamic therapy. *Photochem. Photobiol.* **57**, 242-247.
- Oleinick, N. L. (1998) Apoptosis in response to photodynamic therapy. *Photodynamic News* **1**, 6-9.
- Olson, K. A., J. W. Fett, T. C. French, M. E. Key and B. L. Vallee (1995) Angiogenin antagonists prevent tumor growth *in vivo*. *Proc. Natl. Sci. USA* **92**, 442-446.
- Oseroff, A. R., D. Ohuoha, T. Hasan, J. C. Bommer and M. L. Yarmush (1986) Antibody-targeted photolysis: Selective photodestruction of human T-cell leukemia cells using monoclonal antibody-chlorin e₆ conjugates. *Proc. Natl. Acad. Sci. USA* **83**, 8744-8748.
- Oseroff, A. R., G. Ara, D. Ohuoha, J. Aprille, J. C. Bommer, M. L. Yarmush, J. Foley and L. Cincotta (1987) Strategies for selective cancer photochemotherapy: antibody-targeted and selective carcinoma cell photolysis. *Photochem. Photobiol.* **46**, 83-96.
- Palamanda, J. R., D. Hickman, A. Ward, E. Sim, M. Romkes-Sparks and J. D. Enadkat (1995) Dapsone acetylation by human liver arylamine N-acetyltransferases and interaction with antiopportunistic infection drugs. *Drug Metabolism and Disposition* **23**, 473-477.
- Paquette, B., H. Ali, R. Langlois and J. E. van Lier (1988) Biological Activities of phthalocyanines-VIII. Cellular distribution in V-79 Chinese Hamster Cells and phototoxicity of selectively sulfonated aluminum phthalocyanines. *Photochem. Photobiol.* **47**, 215-220.
- Paquette, B. and J. E. van Lier (1991) Phthalocyanines pour la thérapie photodynamique du cancer: Effet des substituants tertio-butyles sur l'accumulation cellulaire et l'activité photodynamique de phthalocyanines sulfonées de gallium. *J. Chim. Phys.* **88**, 1113-1123.

- Paquette, B. and J. E. van Lier (1991) Phthalocyanines and related compounds: Structure-activity relationships. In *Photodynamic Therapy: Basic principles and clinical applications* (Edited by B. W. Henderson and T. J. Dougherty), pp. 145-156. Marcel Dekker, Inc., New York.
- Parker, J. G. (1987) Optical monitoring of singlet oxygen generation during photodynamic treatment of tumors. *IEEE Circuits and Devices*, 10-21.
- Pasqualini, R. and E. Ruoslahti (1996) Organ targeting *in vivo* using phage display peptide libraries. *Nature* **380**, 364-366.
- Pasqualini, R. E. Koivunen and E. Ruoslahti (1997) αv integrins as receptors for tumor targeting by circulating ligands. *Nature Biotech.* **15**, 542-546.
- Pass, H. I. (1993) Photodynamic therapy in oncology: Mechanisms and clinical use. *J. Natl. Cancer Inst.* **85**, 443-456.
- Patterson, M. S., B. C. Wilson and R. Gaff (1990) *In vivo* tests of the concept of photodynamic threshold dose in normal rat liver photosensitized by aluminum chlorosulphonated phthalocyanine. *Photochem. Photobiol.* **51**, 342-349.
- Payne, C. M., C. Bernstein and H. Bernstein (1995) Apoptosis overview emphasizing the role of oxidative stress, DNA damage and signal transduction pathways. *Leuk. Lymphoma* **19**, 43-93.
- Peeva, M., S. Shopova, N. Stoichkova, N. Michailov, D. Wöhrle and S. Müller (1999) Comparative photodynamic therapy of B16 pigmented melanoma with different generations of sensitizers. *J. Porphyrins and Phthalocyanines* **3**, 380-387.
- Pellegrini, R., S. Martignone, S. Ménard and M. I. Colnaghi (1994) Laminin receptor expression and function in small-cell lung carcinoma. *Int. J. Cancer* **S8**, 116-120.
- Peng, Q., G. W. Farrants, K. Madslien, J. C. Bommer, J. Moan, H. E. Danielsen and J. M. Nesland (1991) Subcellular localization, redistribution and photobleaching of sulfonated aluminum phthalocyanines in a human melanoma cell line. *Int. J. Cancer* **49**, 290-295.
- Peng, Q. and J. Moan (1995) Correlation of distribution of sulphonated aluminium phthalocyanines with their photodynamic effect in tumour and skin of mice bearing CaD2 mammary carcinoma. *Br. J. Cancer* **72**, 565-574.
- Peng, Q., K. Berg, J. Moan, M. Kongshaug and J. M. Nesland (1997) 5-aminolevulinic acid-based photodynamic principles and experimental research. *Photochem. Photobiol.* **65**, 235-251.
- Peng, Q., T. Warloe, K. Berg, J. Moan, M. Kongshaug, K.-E. Giercksky and J. M. Nesland (1997) 5-Aminolevulinic acid-based photodynamic therapy. *Cancer* **79**, 2282-2308.
- Perks, C. M., S. Bowen, Z. P. Gill, P. V. Newcomb and J. M. P. Holly (1999) Differential IGF-independent effects of insulin-like growth factor binding proteins (1-6) on apoptosis of breast epithelial cells. *J. Cellul. Biochem.* **75**, 652-664.
- Peter, K. and T. E. O'Toole (1995) Modulation of cell adhesion by changes in $\alpha_L\beta_2$ (LFA-1, CD11a/CD18) cytoplasmic domain/cytoskeleton interaction. *J. Exp. Med.* **181**, 315-326.
- Peterson, C., M. Masquelier, M. Rudling, K. Söderberg and S. Vitols (1991) Lipoproteins, malignancy and anticancer drugs. In *Lipoproteins as Carriers of Pharmacological Agents* (Edited by J. M. Shaw) New York: Marcel Dekker, pp. 175-200.
- Peterson, C. M., J. M. Lu, Y. Sun, C. A. Peterson, J. G. Shiah, R. C. Straight and J. Kopecek (1996) Combination chemotherapy and photodynamic therapy with N-(2-hydroxypropyl)methacrylamide copolymer-bound anticancer drugs inhibit human ovarian carcinoma heterotransplanted in nude mice. *Cancer Res.* **56**, 3980-3985.

- Pfeiffer, H. (1911) Der nachweis photodynamischer wirkungen fluoreszierender stoffe am lebenden warmblüter. In Handbuch der Biochemischen Arbeitsmethoden (ed. E Abderhalden) vol 1. Berlin: Urban & Schwarzenberg, pp. 563-571.
- PharMingen (1998) Apoptosis, applied reagents and technologies. *Instruction Manual.*, 1st Ed. Becton Dickinson Co.
- Philips, D. (1997) Chemical mechanisms in photodynamic therapy with phthalocyanines. *Progress in Reaction Kinetics* 22, 175-300.
- Pierschbacher, M. D. and E. Ruoslahti (1987) Influence of stereochemistry of the sequence Arg-Gly-Asp-Xaa on binding specificity in cell adhesion. *J. Biol. Chem.* 262, 17294-17298.
- Pierschbacher, M. D., J. W. Polarek, W. S. Craig, J. F. Tschopp, N. J. Sipes and J. R. Harper (1994) Manipulation of cellular interactions with biomaterials toward a therapeutic outcome: a perspective. *J. Cellular Biochem.* 56, 150-154.
- Plow, E. F., M. D. Pierschbacher, E. Ruoslahti, G. A. Marguerie and M. H. Ginsberg (1985) The effect of Arg-Gly-Asp-containing peptides on fibrinogen and von Willebrand factor binding to platelets. *Proc. Natl. Acad. Sci. USA* 82, 8057-8061.
- Polo, L., G. Bianco, E. Reddi and G. Jori (1995) The effect of different liposomal formulations on the interaction of Zn(II)-phthalocyanine with isolated low and high density lipoproteins. *Int. J. Biochem. Cell Biol.* 27, 1249-1255.
- Potts, J. R. and I. D. Campbell (1994) Fibronectin structure and assembly. *Curr. Opinion Cell Biol.* 6, 648-655.
- Povey, A. C. and I. K. O. O'Neill (1990) Copper phthalocyanine labelled magnetic microcapsules: preparation and binding properties *in vitro* and *in vivo* for mutagens having planar molecular structure. *Carcinogenesis* 11, 1989-1993.
- Prosser, E. S., C. Gleeson, K. Carroll and R. O'Kennedy (1991) The effect of immunophotodynamic therapy using the monoclonal antibody HMFG-1 linked to Hematoporphyrin on MCF-7 cell *in vitro*. *Biochem. Soc. Trans.* 20, 53S.
- Putnam, D. and J. Kopecek (1995) Polymer conjugates with anticancer activity. *Adv. Polym. Sci.* 122, 55-123.
- Qin, X. Q., Y. Xiang, C. X. Guan, C. Q. Zhang and X. H. Sun (2001) Integrin-ligands binding reaction upregulates the antioxidant activity of rabbit bronchial epithelial cells. *Sheng Li Hsueh Pao-Acta Physiologica Sinica* 53, 41-44.
- Raab, O. (1900) Über die wirkung fluoreszierender stoffe auf infusoria. *Z. Biol.* 39, 524-526.
- Rahman, S., X. Lu, V. V. Kakkar and K. S. Authi (1995) The integrin $\alpha_{\text{IIb}}\beta_3$ contains distinct and interacting binding sites for snake-venom RGD (Arg-Gly-Asp) proteins. *Biochem. J.* 312, 223-232.
- Rakestraw, S. L., R. G. Tompkins and M. L. Yarmush (1990) Antibody-targeted photolysis: in vitro studies with Sn(IV) chlorin e6 covalently bound to monoclonal antibodies using a modified dextran carrier. *Proc. Natl. Acad. Sci. USA* 87, 4217-4221.
- Rakestraw, S. L., R. G. Tompkins and M. L. Yarmush (1990) Preparation and characterization of immunoconjugates for antibody-targeted photolysis. *Bioconjug. Chem.* 1, 212-221.
- Rakestraw, S. L., W. E. Ford, R. G. Tompkins, M. A. Rodgers, W. P. Thorpe and M. L. Yarmush (1992) Antibody-targeted photolysis: in vitro immunological, photophysical and cytotoxic properties of monoclonal antibody-dextran-Sn(IV) chlorin e6 immunoconjugates. *Biotech. Prog.* 8, 30-39.

- Reddi, E., C. Zhou, R. Biolo, E. Menegaldo, G. Jori (1990) Liposome- or LDL-administered Zn(II)-phthalocyanine as a photodynamic agent for tumours. I. Pharmacokinetic properties and phototherapeutic efficiency. *Br. J. Cancer* **61**, 407-411.
- Reddi, E. (1997) Role of delivery vehicles for photosensitizers in the photodynamic therapy of tumours. *J. Photochem. Photobiol. B: Biol.* **37**, 189-195.
- Redman, C., M. J. Xu, Y.-M. Peng, J. A. Scott, C. Payne, L. C. Clarke and M. A. Nelson (1997) Involvement of polyamines in selenomethionine induced apoptosis and mitotic alterations in human tumor cells. *Carcinogenesis* **18**, 1195-1202.
- Redmond, R. W. (1991) Enhancement of the sensitivity of radiative and non-radiative detection techniques in the study of photosensitization by water-soluble sensitizers using a reverse micelle system. *Photochem. Photobiol.* **54**, 547-556.
- Rembaum, A. and Z. A. Tokés (1989) *Microspheres: Medical and Biological Applications*. CRC Press, Boca Raton.
- Rensen, P. C. N., W. G. Love and P. W. Taylor (1994) In vitro interaction of zinc(II)-phthalocyanine-containing liposomes and plasma proteins. *J. Photochem. Photobiol. B: Biol.* **26**, 29-35.
- Ribova, B., N. L. Krinick and J. Kopecek (1993) Targetable photoactivatable drugs 3. In vitro efficacy of polymer bound chlorin-e6 toward human hepatocarcinoma cell line (PLC/PRF/5) targeted with galactosamine and to mouse splenocytes targeted with anti-Thy 1.2 antibodies. *J. Controlled Rel.* **25**, 71-87.
- Ricchelli, F. (1995) Photophysical properties of porphyrins in biological membranes. *J. Photochem. Photobiol. B: Biol.* **29**, 109-118.
- Richter, A. M., B. Kelly, J. Chow, D. J. Liu, G. H. N. Towers, D. Dolphin and J. G. Levy (1987) Preliminary studies on a more effective phototoxic agent than hematoporphyrin. *JNCI*. **79**, 1327-1332.
- Richter, A. M., E. Cerruti-Sola, E. D. Sternberg, D. Dolphin and J. G. Levy (1990) Biodistribution of tritiated benzoporphyrin derivative (³H-BPD-MA), a new potent photosensitizer, in normal and tumor-bearing mice. *J. Photochem. Photobiol. B: Biol.* **5**, 231-244.
- Richter, A. M., E. Waterfield, A. K. Jain, A. J. Canaan, B. A. Allison and J. G. Levy (1993) Liposomal delivery of a photosensitizer benzoporphyrin derivative monoacid ring A (BPD), to tumor tissue in a mouse tumor model. *Photochem. Photobiol.* **57**, 1000-1006.
- Ris, H. B., T. Krueger, A. Giger, C. K. Lim, J. C. Stewart, U. Althaus and H. J. Altermatt (1999) Photodynamic therapy with mTHPC and polyethylene glycol-derived mTHPC: a comparative study on human tumour xenografts. *Br. J. Cancer* **79**, 1061-1066.
- Rivellese, M. J. and C. R. Bauman (1999) Photodynamic therapy of eye diseases. *Ophthalm. Surg. Lasers* **30**, 653-661.
- Robert, R. J., K. E. O'Neill and C. E. Yen (1984) DNA sequences from the adenovirus 2 genome *J. Biol. Chem.* **259**, 13968-13975.
- Robertson, J. M. (1935) An X-ray study of the structure of the phthalocyanines. Part I. The metal-free, nickel, copper and platinum compounds. *J. Chem. Soc.* 615-621.
- Robertson, J. M. (1936) An X-ray study of the structure of the phthalocyanines. Part II. Quantitative structure determination of the metal-free compound. *J. Chem. Soc.* 1195-1209.
- Robertson, J. M. and I. Woodward (1937) An X-ray study of the phthalocyanines. Part III. Quantitative structure determination of nickel phthalocyanine. *J. Chem. Soc.* 219-230.
- Robertson, J. M. and I. Woodward (1940) An X-ray study of the phthalocyanines. Part IV. Direct quantitative analysis of the platinum compound. *J. Chem. Soc.* 36-48.

- Röder, B., S. Hackbarth and A. Paul (2000) Photophysical studies on pheophorbide a – a promising sensitizer for PDT: Ways to increase the selectivity and the local concentration in target cells. 13th International Congress on Photobiology July 1-6, 2000; San Francisco, California: Abstract 561.
- Rosenkranz, A. A., D. A. Jans and A. S. Sobolev (2000) Targeted intracellular delivery of photosensitizers to enhance photodynamic therapy. *Immunol. Cell Biol.* 78, 452-464.
- Rosenthal, I. (1991) Phthalocyanines as photodynamic sensitizers. *Photochem. Photobiol.* 53, 859-870.
- Rosenthal, D. I. and E. Glatstein (1994) Clinical applications of photodynamic therapy. *Ann. Med.* 26, 405-409.
- Rosenthal, I. and E. Ben-Hur (1995) Role of oxygen in the phototoxicity of phthalocyanines. *Int. J. Radiat. Biol.* 67, 85-91.
- Ross, A. H. and H. M. McConnell (1975) Permeation of a spin-labeled phosphate into the human erythrocyte. *Biochem.* 14, 2793-2803.
- Rovati, G. E. (1993) Rational experimental design and data analysis for ligand binding studies tricks, tips and pitfalls. *Pharmacological Res.* 28, 277-299.
- Ruebner, A., D. Kirsch, S. Andrees, W. Decker, B. Roeder, B. Spengler, R. Kaufman and J. G. Moser (1997) Dimeric cyclodextrin carriers with high binding affinity to porphyrinoid photosensitizers. *J. Inclusion Phenom.* 27, 69-84.
- Ruebner, A., Z. Yang, D. Leung and R. Breslow (1999) A cyclodextrin dimer with a photocleavable linker as a possible carrier for the photosensitizer in photodynamic tumor therapy. *Proc. Natl. Acad. Sci. USA* 96, 14692-14693.
- Ruolahti, E. (1996) RGD and other recognition sequences for integrins. *Annu. Rev. Cell Biol.* 12, 697-715.
- Ruolahti, E. (1997) Integrins as signaling molecules and targets for tumor therapy. *** *International* 51, 1413-1417.
- Ruolahti, E. and A. Vaheri (1997) Cell-to-cell contact and extracellular matrix. *Curr. Opinion Cell Biol.* 9, 605-607.
- Ruolahti, E. (1999) Fibronectin and its integrin receptors in cancer. *Adv. Cancer Res.* 1-20.
- Ruolahti, E. and J. Reed (1999) New way to activate caspases. *Nature* 397, 479-480.
- Ruolahti, E. (2000) Targeting tumor vasculature with homing peptides from phage display. *Sem Cancer Biol.* 10, 435-442.
- Ruolahti, E. and D. Rajotte (2000) An address system in the vasculature of normal tissues and tumors. *Annu. Rev. Immunol.* 18, 813-827.
- Saiki, I., J. Iida, J. Murata, R. Ogawa, N. Nishi, K. Sugimura, S. Tokura and I. Azuma (1989) Inhibition of the metastasis of murine malignant melanoma by synthetic polymeric peptides containing core sequences of cell-adhesive molecules. *Cancer Res.* 49, 3815-3822.
- Santos, A. E., J. A. N. Laranjinha and L. M. Almeida (1998) Sulfonated chloroaluminum phthalocyanine incorporates into human plasma lipoproteins: Photooxidation of low-density lipoproteins. *Photochem. Photobiol.* 67, 378-385.
- Sastry, S. K. and A. F. Horwitz (1993) Integrin cytoplasmic domains: mediators of cytoskeletal linkages and extra- and intracellular initiated transmembrane signaling. *Curr. Opinion Cell Biol.* 5, 819-831.
- Schafer, A. I. (1996) Antiplatelet therapy. *Am. J. Med.* 101, 199-209.
- Schmidt, S., U. Wagner, B. Schultes, P. Oehr, W. Decleer, W. Ertmer, H. Lubaschowski, H. J. Biersack and D. Krebs (1992) [Photodynamic laser therapy with antibody-bound dyes. A new procedure in therapy of gynecologic malignancies (German)]. *Fortschritte der Medizin* 110, 298-301.

- Schmidt-Erfurth, U., W. Bauman, E. Gragoudas, T. J. Flotte, N. A. Michaud, R. Birngruber and T. Hasan (1994) Photodynamic therapy of experimental choroidal melanoma using lipoprotein-delivered benzoporphyrin. *Ophthalmology* **101**, 89-99.
- Schmidt-Erfurth, U., T. J. Flotte, E. Gragoudas, K. Schomacker, R. Birngruber and T. Hasan (1996) Benzoporphyrin-lipoprotein-mediated photodestruction of intraocular tumors. *Exp. Eye Res.* **62**, 1-10.
- Schmidt-Erfurth, U., Diddens H, Birngruber R, Hasan T.(1997) Photodynamic targeting of human retinoblastoma cells using covalent low-density lipoprotein conjugates. *Br J Cancer* **75**, 54-61.
- Schuitmaker, J. J., P. Baas, H. L. L. M. van Leengoed, F. W. van der Meulen, W. M. Star and N. van Zandwijk (1996) Photodynamic therapy: a promising new modality for the treatment of cancer. *J. Photochem. Photobiol. B: Biol.* **34**, 3-12.
- Schuller, H. M., K. L. Becker and H.-P. Witschi (1988) An animal model for neurocrine lung cancer. *Carcinogenesis* **9**, 293-296.
- Schuster, H., B. Rautenstrauss, M. Mittag, D. Stratmann and E. Schweizer (1995) Substrate and product binding sites of yeast fatty acid synthase, Stoichiometry and binding kinetics of wild-type and *in vitro* mutated enzymes. *Eur. J. Biochem.* **228**, 417-424.
- Scott, G., L. Cassidy, and A. Busacco (1997) Fibronectin suppresses apoptosis in normal human melanocytes through an integrin-dependent mechanism. *J. Invest. Dermatol.* **108**, 147-153.
- Segalla, A., C. Milanese, G. Jori, H.-G. Capraro, U. Isele and K. Schiesek (1994) CGP 55398, a liposomal Ge(IV) phthalocyanine bearing two axially ligated cholesterol moieties: a new potential agent for photodynamic therapy of tumours. *Br. J. Cancer* **69**, 817-825.
- Selbo, P. K., G. Sivam, Ø Fodstad, K. Sandvig and K. Berg (2000) Photochemical internalisation increases the cytotoxic effect of the immunotoxin MOC31-gelonin. *Int. J. Cancer* **87**, 853-859.
- Selman, S. H., G. M. Garbo, R. W. Keck, M. Kreimer-Birnbaum and A. R. Morgan (1987) A dose response analysis of purpurin derivatives used as photosensitizers for the photodynamic treatment of transplantable FANFT induced urothelial tumors. *J. Urol.* **137**, 1255-1257.
- Separovic, D., J. He and N. L. Oleinick (1997) Ceramide generation in response to photodynamic treatment of L5178Y mouse lymphoma cells. *Cancer Res.* **57**, 1717-1721.
- Separovic, D., K. J. Mann and N. L. Oleinick (1998) Association of ceramide accumulation with photodynamic treatment-induced cell death. *Photochem. Photobiol.* **68**, 101-109.
- Separovic, D. K., J. J. Pink, N. A. Oleinick, M. Kester, D. A. Boothman, M. McLoughlin, L. A. Peña and A. Haimovitz-Friedman (1999) Niemann-Pick human lymphoblasts are resistant to phthalocyanine 4-photodynamic therapy-induced apoptosis. *Biochem. Biophys. Res. Commun.* **258**, 506-512.
- Sethi, T., R. C. Rintoul, S. M. Moore, A. C. MacKinnon, D. Salter, C. Choo, E. R. Chilvers, I. Dransfield, S. C. Donnelly, R. Strieter and C. Haslett (1999) Extracellular matrix proteins protect small cell lung cancer cells against apoptosis: A mechanism for small cell lung cancer growth and drug resistance *in vivo*. *Nature Med.* **5**, 662-668.
- Seufferlein, T. and E. Rozengurt (1994) Lysophosphatidic acid stimulates tyrosine phosphorylation of focal adhesion kinase, paxillin and p130. *J. Biol. Chem.* **269**, 9345-9351.
- Sharman, W. M., C. M. Allen and J. E. van Lier (1999) Photodynamic therapeutics: Basic principles and clinical applications. *Drug Discovery Today* **4**, 507-517.
- Sharman, W. M., C. M. Allen and J. E. van Lier (2000) Role of activated oxygen species in photodynamic therapy. *Methods Enzymol.* **319**, 376-400.
- Sheppard, D. (1996) Epithelial integrins. *BioEssays* **18**, 655-660.
- Shiah, J.-G., Y. Sun, M. Peterson and J. Kopecek (1999) Biodistribution of free and N-(2-hydroxypropyl)methacrylamide copolymer-bound mesochlorin e₆ and adriamycin in nude mice bearing human ovarian carcinoma OVCAR-3 xenografts. *J. Controlled Release* **61**, 145-157.

- Shiah, J. G., Y. Sun, C. M. Peterson, R. C. Straight and J. Kopecek (2000) Antitumor activity of N-(2-hydroxypropyl)methacrylamide copolymer-mesochlorine e6 and adriamycin conjugates in combination treatments. *Clin. Cancer Res.* **6**, 1008-1015.
- Shirk, J. S. (1996) Nonlinear optical properties of metallophthalocyanines. in *Phthalocyanines: Properties and Applications Volume 1* (eds. C. C. Leznoff and A. B. P. Lever). New York: VCH Publishers, Inc., pp. 79-181.
- Shyy, A. J-Y and S. Chien (1997) Role of integrins in cellular responses to mechanical stress and adhesion. *Curr. Opinion Cell Biol.* **9**, 707-713.
- Sickenberg, M., U. Schmidt-Erfurth, J. W. Miller, C. J. Pournaras, L. Zografos, B. Piguet, G. Donati, H. Laqua, I. Barbazetto, E. S. Gragoudas, A.-M. Lane, R. Birngruber, H. van den Bergh, A. H. Strong, U. Manjuris, T. Gray, M. Fasdn and N. M. Bressler (2000) A preliminary study of photodynamic therapy using verteporfin for choroidal neovascularization in pathologic myopia, ocular histoplasmosis syndrome, angioid streaks, and idiopathic causes. *Arch. Ophthalmol.* **118**, 327-336.
- Singh, A. (1982) Chemical and biochemical aspects of superoxide radicals and related species of activated oxygen. *Can. J. Physiol. Pharmacol.* **60**, 1330-1345.
- Sinn, H., H. H. Schrenk, E. A. Friedrich, U. Schilling and W. Maier-Borst (1990) Design of compounds having an enhanced tumour uptake, using serum albumin as a carrier. Part I. *Nucl. Med. Biol.* **17**, 819-827.
- Skalkos, D., J. A. Hampton, R. W. Keck, M. Wagoner and S. H. Selman (1994) Iminium salt benzochlorins: structure-activity relationship studies. *Photochem. Photobiol.* **59**, 175-181.
- Skorobogaty, A., T. D. Smith, G. Dougherty and J. R. Pilbrow (1985) Synthesis and physico-chemical properties of cationic derivatives of phthalocyaninatocopper (II). *J. Chem. Soc. Dalton Trans.* 651-658.
- Snow, A. W. and W. R. Barger (1989) Phthalocyanine films in chemical sensors. in *Phthalocyanines: Properties and Applications Volume 1* (eds. C. C. Leznoff and A. B. P. Lever). New York: VCH Publishers, Inc., pp. 341-392.
- Sobolev, A. S. and E. F. Stranadko (1997) Photodynamic therapy in Russia: Clinical and fundamental aspects. *International Photodynamics* **1**, 2-3.
- Sobolev, A. S., D. A. Jans and A. A. Rosenkranz (2000) Targeted intracellular delivery of photosensitizers. *Prog. Biophys. Mol. Biol.* **73**, 51-90.
- Sokolov, V. V., V. I. Chissov, E. V. Filonenko, G. M. Sukhin, R. I. Yakubovskaya, T. A. Belous, N. N. Zharkova, D. N. Kozlov and V. V. Smirnov (1994) Photodynamic therapy of cancer with photosensitizer PHOTOGEN. *Proc. SPIE* **2325**, 367-374.
- Sokolov, V. V., V. I. Chissov, E. V. Filonenko, R. I. Yakubovskaya, D. G. Sukhin, M. G. Galpern, G. N. Vorozhtsov, A. V. Gulin, M. B. Zhitkova, N. N. Zharkova, D. N. Kozlov and V. V. Smirnov (1994) First clinical results with new drug for PDT. *Proc. SPIE* **2325**, 364-366.
- Soncin, M., L. Polo, E. Reddi, G. Jori, M. E. Kenney, G. Cheng and M. A. J. Rodgers (1995) Effect of axial ligation and delivery system on the tumour-localising and -photosensitising properties of Ge(IV)-octabutoxyphthalocyanine. *Br. J. Cancer* **71**, 727-732.
- Soukos, N. S., M. R. Hamblin and T. Hasan (1997) The effect of charge on cellular uptake and phototoxicity of polylysine chlorin e6 conjugates. *Photochem. Photobiol.* **65**, 723-729.
- Spikes, J. D. (1992) Quantum yields and kinetics of the photobleaching of hematoporphyrin, photofrinII, tetra(4-sulfonatophenyl)-porphine and uroporphyrin. *Photochem. Photobiol.* **55**, 797-808.
- Stables, G. I. (1999) Photodynamic therapy in dermatology. *J. Dermatol. Treat.* **10**, 213-219.
- Steck, T. L., G. Fairbanks and D. F. H. Wallach (1971) Disposition of the major proteins in the isolated erythrocyte membrane, proteolytic dissection. *Biochem.* **10**, 2617-2624.

- Stein, R., R. Blumenthal, R. M. Sharkey and D. M. Goldenberg (1994) Comparative biodistribution and radioimmunotherapy of monoclonal antibody RS7 and its F(ab')₂ in nude mice bearing human tumor xenografts. *Cancer* **73**, 816-823.
- Stewart, M. M. Thiel and N. Hogg (1995) Leukocyte integrins. *Curr. Opinion Cell Biol.* **7**, 690-696.
- Stoner, G. D., G. Adam-Rodwell and M. A. Morse (1993) Lung tumors in strain A mice: Application for studies in cancer chemoprevention. *J. Cellular Biochem.* **S17F**, 95-103.
- Strandadko, E. F., O. K. Skobelkin, G. D. Litvin and T. A. Astrakhankina (1996) Photodynamic therapy of human malignant tumours: a comparative study between photohem and tetrasulfonated aluminium phthalocyanine. *Proc. SPIE* **2728**, 194-202.
- Strömblad, S. and D. A. Cheresh (1996) Integrins, angiogenesis and vascular cell survival. *Chem. & Biol.* **3**, 881-885.
- Strömblad, S., J. C. Becker, M. Yebra, P. C. Brooks and D. A. Cheresh (1996) Suppression of p53 activity and p21^{WAF1/CIP1} expression by vascular cell integrin $\alpha v\beta 3$ during angiogenesis. *J. Clin. Invest.* **98**, 426-433.
- Strong, L., D. M. Yarmush and M. L. Yarmush (1994) Antibody-targeted photolysis. *Annal. NY Acad. Sci.* **745**, 297-320.
- Subramanian, T., B. Tarodi and G. Chinnadurai (1995) p53-independent apoptotic and necrotic cell deaths induced by adenovirus infection: Suppression by E1B 19K and Bcl-2. *Cell Growth and Differentiation*. **6**, 131-137.
- Sugiyama, T., S. Kumagai, T. Nishida, K. Ushijima, T. Matsuo, M. Yakushiji, S. H. Hyon and Y. Ikada (1998) Experimental and clinical evaluation of cisplatin-containing microspheres as intraperitoneal chemotherapy for ovarian cancer. *Anticancer Res.* **18**, 2837-2842.
- Sundaram, S. G. and J. A. Milner (1996) Diallyl disulfide induces apoptosis of human colon tumor cells. *Carcinogenesis* **17**, 669-673.
- Sutton, J. M., N. Fernandez and R. W. Boyle (2000) Functionalized diphenylchlorins and bacteriochlorins: Their synthesis and bioconjugation for targeted photodynamic therapy and tumour cell imaging. *J. Porphyrins Phthalocyanines* **4**, 655-658.
- Suzuki, S., Ta. Takahashi, S. Nakamura, K. Koike, Y. Ariyoshi, To. Takahashi and R. Ueda (1993) Alterations in integrin expression in human lung cancer. *Jpn. J. Cancer Res.* **84**:168-174.
- Szejtli, J. (1994) Medicinal applications of cyclodextrins. *Med. Research Rev.* **14**, 353-386.
- Tada, H., O. Shiho, K. Kuroshima, M. Koyama and K. Tsukamoto (1986) An improved colorimetric assay for interleukin 2. *Journal of Immunological Methods* **93**, 157-165.
- Taillefer, J., M.-C. Jones, N. Brasseur, J. E. van Lier and J.-C. Leroux (2000) Preparation and characterization of pH-responsive polymeric micelles for the delivery of photosensitizing anticancer drugs. *J. Pharma. Sci.* **89**, 52-62.
- Takeichi, M. (1993) Cadherins in cancer: implications for invasion and metastasis. *Curr. Opinion Cell Biol.* **5**, 806-811.
- Takemura, T., N. Ohta, S. Nakajima and I. Sakata (1989) Critical importance of the triplet lifetime of the photosensitizer in photodynamic therapy of tumors. *Photochem. Photobiol.* **50**, 339-344.
- Tappeiner, H. V. and A. Jesionek (1903) Therapeutische versuche mit fluoreszierenden stoffe. *Muench. Med. Wochenschr.* **47**, 2042-2044.
- Thomas, J. P. and A. W. Girotti (1988) Photogeneration of singlet oxygen by membrane-bound hematoporphyrin derivative. *Photochem. Photobiol.* **47**, 47S.
- Thompson, D. H., O. V. Gerasimov, J. J. Wheeler, Y. Rui and V. C. Aderson (1996) Triggerable plasmalogen liposomes: improvement of system efficiency. *Biochimica Biophysica Acta* **1279**, 25-34.

- Thornberry, N. A. and Y. Lazebnik (1998) Caspases: The enemies within. *Science* **281**, 1312-1316.
- Uhal, B. D., I. Joshi, A. L. True, S. Mundle, A. Raza, A. Pardo and M. Selman (1995) Fibroblasts isolated after fibrotic lung injury induce apoptosis of alveolar epithelial cells in vitro. *Am. J. Physiol.* **269**, L819-L828.
- Urizzi, P., J.-P. Souchard, C. Palevody, G. Ratovo, E. Hollande and F. Nepveu (1997) LDL labeled with indium-111 through a lipid anchor for tumor localization: internalization in human pancreatic cancer cells maintained in culture. *Int. J. Cancer* **70**, 1-8.
- Urizzi, P., C. M. Allen, R. Langlois, R. Ouellet, C. La Madeleine and J. E. van Lier (2001) Low-density lipoprotein-bound aluminum sulfophthalocyanine: Targeting tumor cells for photodynamic therapy. *J. Porphyrins Phthalocyanines* **5**, 154-160.
- Vakoulovskaja, E. G. V. V. Chental, N. A. Abdoullin, Y. P. Kuvshinov, T. D. Tabolinovskaja, N. J. Edinak, B. K. Poddubny, V. L. Lioubaev, V. P. Boikov, T. T. Kondratjeva, G. A. Meerovich, A. A. Stratonnikov, K. G. Linkov and V. V. Agafonov (1994) Photodynamic therapy of head and neck tumors. *Proc SPIE* **2924**, 309-312.
- Valduga, G., G. Bianco, G. Csik, E. Reddi, L. Masiero, S. Garbisa and G. Jori (1996) Interaction of hydro- or lipophilic phthalocyanines with cells of different metastatic potential. *Biochem. Pharmacol.* **51**, 585-590.
- Valduga, G., E. Reddi, S. Garbisa and G. Jori (1998) Photosensitization of cells with different metastatic potential by liposome-delivered Zn(II)-phthalocyanine. *Int. J. Cancer* **75**, 412-417.
- Valduga, G., A. Pifferi, G. Taroni, E. Reddi, R. Cubeddu and G. Jori (1999) Steady-state and time resolved spectroscopic studies on low density lipoprotein bound Zn(II)Pc. *J. Photochem. Photobiol. B: Biol.* **49**, 198-203.
- Valenzano, D. P. (1987) Photomodification of biological membranes with emphasis on singlet oxygen mechanisms. *Photochem. Photobiol.* **46**, 147-160.
- van Hillegersberg, W. J. Kort and J. H. P. Wilson (1994) Current status of photodynamic therapy in oncology. *Drugs* **48**, 510-527.
- van Hillegersberg, R., J. M. Hekking-Weijma, J. H. P. Wilson, A. Edixhoven-Bosdijk and W. J. Kort. (1995) Adjuvant intraoperative photodynamic therapy diminishes the rate of local recurrence in a rat mammary tumour model. *Br. J. Cancer* **71**, 733-737.
- van Leengoed, H. L. L. M., V. Cuomo, A. A. C. Versteeg, N. van der Veen, G. Jori, W. M. Star (1994) *In vivo* fluorescence and photodynamic activity of zinc phthalocyanine administered in liposomes. *Br. J. Cancer* **69**, 840-845.
- van Lier, J. E. (1988) New sensitizers for photodynamic therapy of cancer. In *Light in Biology and Medicine* Vol 1 (eds Douglas, R. H., J. Moan and F. Dell'Acqua) New York: Plenum Publishing, pp. 133-140.
- van Lier, J. E., N. Brasseur, B. Paquette, J. R. Wagner, H. Ali, R. Langlois and J. Rousseau (1988) Phthalocyanines as sensitizers for photodynamic therapy of cancer. In *Photosensitization* (ed. Moreno, G.) Berlin: Springer-Verlag, pp. 435-444.
- van Lier, J. E. and J. D. Spikes (1989) The chemistry, photophysics and photosensitizing properties of phthalocyanines. in *Photosensitizing compounds: Their chemistry, biology and clinical use*. Chichester: John Wiley and Sons, Ltd., pp. 17-32.
- van Lier, J. E. (1990) Phthalocyanines as sensitizers for the PDT of cancer. In *Photodynamic therapy of neoplastic disease* (Edited by D. Kessel), pp. 279-291. CRC Press, Boca Raton.
- van Peteghem, M. C., M. M. Marcel and G. K. de Bruyne (1980) Phagocytic capacity of invasive malignant cells in three-dimensional culture. *Virchows Arch. B. Cell Pathol.* **193**, 193-204.
- Varner, J. A. and D. A. Cheresh (1996) Integrins and cancer. *Curr. Opinion in Cell Biol.* **8**, 724-730.

- Varnes, M. E., M. T. Bayne and G. R. Bright (1996) Reduction of intracellular pH is not the mechanism for the synergistic interaction between photodynamic therapy and nigericin. *Photochem. Photobiol.* **64**, 853-858.
- Varnes, M. E., S.-M. Chiu, L.-Y. Xue and N. L. Oleinick (1999) Photodynamic therapy-induced apoptosis in lymphoma cells: translocation of cytochrome c causes inhibition of respiration as well as caspase activation. *Biochem. Biophys. Res. Comm.* **255**, 673-679.
- Versluis, A. J., P. C. N. Rensen, M. E. Kuipers, W. G. Love and P. W. Taylor (1994) Interaction between zinc(II)-phthalocyanine-containing liposomes and human low density lipoproteins. *J. Photochem. Photobiol. B: Biol.* **23**, 141-148.
- Viola, A., A. Jeunet, R. Decreau, M. Chanon and M. Julliard (1998) ESR studies of a series of phthalocyanines. Mechanism of phototoxicity. Comparative quantitation of $O_2^{\cdot -}$ using ESR spin-trapping and cytochrome c reduction techniques. *Free Rad. Res.* **28**, 517-532.
- Vitols, S. (1991) Uptake of low-density lipoprotein by malignant cells. Possible therapeutic applications. *Cancer Cells* **3**, 488-495.
- von Tappeiner, H. and A. Jodlbauer (1907) Die sensibilisierende wirkung fluorescienrender substanzen: gesammelte untersuchungen über die photodynamische erscheilung. In Vogel FGW (ed) Leipzig, pp. 210.
- Vowels, B. R., E. K. Yoo and F. P. Gasparro (1996) Kinetic analysis of apoptosis induction in human cell lines by UVA and 8-MOP. *Photochem. Photobiol.* **63**, 572-576.
- Vrouenraets, M. B., G. W. M. Visser, F. A. Stewart, M. Stigter, H. Oppelaar, P. E. Postmus, G. B. Snow and G. A. M. S. van Dongen (1999) Development of *meta*-tetrahydroxyphenylchlorin-monoclonal antibody conjugates for photoimmunotherapy. *Cancer Res* **59**, 1505-1513.
- Vrouenraets, M. B., G. W. M. Visser, C. Loup, B. Meunier, M. Stigter, H. Oppelaar, F. A. Stewart, G. B. Snow and G. A. M. S. van Dongen (2000) Targeting of a hydrophilic photosensitizer by use of internalizing monoclonal antibodies: A new possibility for use in photodynamic therapy. *Int. J. Cancer* **88**, 108-114.
- Vuori, K. and E. Ruoslahti (1994) Association of insulin receptor substrate-1 with integrins. *Science* **266**, 1576-1578.
- Wagner, J. R., H. Ali, R. Langlois, N. Brasseur and J. E. van Lier (1987) Biological activities of phthalocyanines VI. Photooxidation of L-tryptophan by selectively sulfonated gallium phthalocyanines: Singlet oxygen yields and effect of aggregation. *Photochem. Photobiol.* **45**, 587-594.
- Wang, H., M. Rubin, E. Fenig, A. DeBlasio, J. Mendelsohn, J. Yahalom and R. Wieder (1997) Basic fibroblast growth factor causes growth arrest in MCF-7 human breast cancer cells while inducing both mitogenic and inhibitory G_1 events. *Cancer Res.* **57**, 1750-1757.
- Weber, J. H. and D. H. Busch (1965) Complexes derived from strong field ligands. XIX. Magnetic properties of transition metal derivatives of 4,4',4'',4'''-tetrasulfophthalocyanines. *Inorg. Chem.* **4**, 469-471.
- Weber, J. M., B. G. Talbot and L. Delorme (1989) The orientation of the adenovirus fiber and its anchor domain identified through molecular mimicry. *Virology* **168**, 180-182.
- Weishaupt, K. R., C. J. Gomer and T. J. Dougherty (1976) Identification of singlet oxygen as the cytotoxic agent in photo-inactivation of a murine tumor. *Cancer Res.* **36**, 2326-2329.
- Westerman, P., T. Glanzmann, S. Andrejevic, D. R. Braichotte, M. Forrer, G. A. Wagnieres, P. Monnier, H. van den Bergh, J. P. Mach and S. Folli (1998) Long circulating half-life and high tumor selectivity of the photosensitizer *meta*-tetrahydroxyphenylchlorin conjugated to polyethylene glycol in nude mice grafted with a human colon carcinoma. *Int. J. Cancer* **76**, 842-850.
- Wheldon, T. E. (1994) Targeting radiation to tumours. *Int. J. Radiat. Biol.* **65**, 109-116.

- Whitlock, B. B., S. Gardai, V. Fadok, D. Bratton and P. M. Henson (2000) Differential roles of $\alpha_M\beta_2$ integrin clustering or activation in the control of apoptosis via regulation of AKT and ERK survival mechanisms. *J. Cell Biol.* **151**, 1305-1320.
- Wickham, T. J., P. Mathias, D. A. Cheresch and G. R. Nemerow (1993) Integrins $\alpha_v\beta_3$ and $\alpha_v\beta_5$ promote adenovirus internalization but not virus attachment. *Cell* **73**, 309-319.
- Wickham, T. J., E. J. Filardo, D. A. Cheresch and G. R. Nemerow (1994) Integrin $\alpha_v\beta_5$ selectively promotes adenovirus mediated cell membrane permeabilization. *J. Cell Biol.* **127**, 257-264.
- Wilson, J. M. (1995) Gene therapy for cystic fibrosis: Challenges and future directions. *J. Clin. Invest.* **96**, 2547-2554.
- Witjes, M. J. H., O. C. Speelman, P. G. J. Nikkels, C. A. A. M. Nooren, B. van der Holt, H. L. L. M. van Leengoed, W. M. Star and J. L. N. Roodenburg (1996) *In vivo* fluorescence kinetics and localisation of aluminium phthalocyanine disulphonate in an autologous tumour model. *Br. J. Cancer* **73**, 573-580.
- Wöhrle, D. (1989) Phthalocyanines in polymer phases. in *Phthalocyanines: Properties and Applications Volume 1* (eds. C. C. Leznoff and A. B. P. Lever). New York: VCH Publishers, Inc., pp. 55-132.
- Wöhrle, D., A. Hirth, T. Bogdahn-Rai, G. Schnurpfeil and M. Shopova (1998) Photodynamic therapy of cancer: Second and third generations of photosensitizers. *Russ. Chem. Bull.* **47**, 807-816.
- Wöhrle, D., S. Müller, M. Shopova, V. Mantareva, G. Spassova, F. Vietri, F. Ricchelli and G. Jori (1999) Effect of delivery system on the pharmacokinetic and phototherapeutic properties of bis(methyloxyethyleneoxy) silicon-phthalocyanine in tumor-bearing mice. *J. Photochem. Photobiol. B: Biol.* **50**, 124-128.
- Woodburn, K., S. Styli, J. S. Hill, A. H. Kaye, J. A. Reiss and D. R. Phillips (1992) Evaluation of tumour and tissue distribution of porphyrins for use in photodynamic therapy. *Br. J. Cancer* **65**, 321-328.
- Woodburn, K. and D. Kessel (1994) The alteration of plasma lipoproteins by Cremophor EL. *J. Photochem. Photobiol. B: Biol.* **22**, 197-201.
- Woodburn, K., C. K. Chang, S. Lee, B. Henderson and D. Kessel (1994) Biodistribution and PDT efficacy of a ketochlorin photosensitizer as a function of the delivery vehicle. *Photochem. Photobiol.* **60**, 154-159.
- Woodburn, K., E. Sykes and D. Kessel (1995) Interactions of Solutol HS 15 and Cremophor EL with plasma lipoproteins. *Int. J. Biochem. Cell Biol.* **27**, 693-699.
- Woodburn, K. W., Q. Fan, D. R. Miles, D. Kessel, Y. Luo and S. W. Young (1997) Localization and efficacy analysis of the phototherapeutic lutetium texaphyrin (PCI-0123) in the murine EMT6 sarcoma model. *Photochem. Photobiol.* **65**, 410-415.
- Woods, S. R., J. A. Holroyd and S. B. Brown (1997) The subcellular localization of Zn(II)phthalocyanines and their redistribution on exposure to light. *Photochem. Photobiol.* **65**, 397-402.
- Wustrow, T. P. U. (1995) Mechanisms of phototoxic effect in photodynamic therapy. *Lasers in Otorhinolaryngology, and in Head and Neck Surgery* **49**, 31-35.
- Xiao, Y. and G. A. Truskey (1996) Effect of receptor-ligand affinity on the strength of endothelial cell adhesion. *Biophysical Journal* **71**, 2869-2884.
- Xu, S. X., N. Q. Wang and Z. W. Dong (1989) [Experimental study of anti-tumor effect with conjugates of monoclonal antibodies and hematoporphyrin derivative (Chinese)]. *Yao Hsueh Hsueh Pao* **24**, 401-406.
- Xue, L.-Y., J. He and N. L. Oleinick (1997) Rapid tyrosine phosphorylation of HS1 in the response of mouse lymphoma L5178Y-R cells to photodynamic treatment sensitized by the phthalocyanine Pc4. *Photochem. Photobiol.* **66**, 105-113.
- Xue, L.-Y., J. He and N. L. Oleinick (1999) Promotion of photodynamic therapy-induced apoptosis by stress kinases. *Cell Death and Differentiation* **6**, 855-864.

- Yamada, K. M. and S. Miyamoto (1995) Integrin transmembrane signaling and cytoskeletal control. *Curr. Opin Cell Biol.* **7**, 681-689.
- Yang, Y., Q. Li, H. C. J. Ertl and J. M. Wilson (1995) Cellular and humoral immune response to viral antigens create barriers to lung-directed gene therapy with recombinant adenoviruses. *J. Virol.* **69**, 2004-2015.
- Yates, N. C. , J. Moan and A. Western (1990) Water-soluble metal naphthalocyanines-near-IR photosensitizers: cellular uptake, toxicity and photosensitizing properties in NHIK 3025 human cancer cells. *J. Photochem. Photobiol. B: Biol.* **4**, 379-390.
- Young, L. H. Y., M. A. Howard, L. K. Hu, R. Y. Kim and E. S. Gragoudas (1996) Photodynamic therapy of pigmented choroidal melanomas using a liposomal preparation of benzoporphyrin derivative. *Arch. Ophthalmol.* **114**, 186-192.
- Zaidi, S. I. A., J. Faulk, B. D. Rihter, M. E. Kenney and H. Mukhtar (1992) Tissue uptake, distribution and photodynamic effects of silicon phthalocyanine administered in corn oil and liposomes to mice bearing transplantable tumors. *Photochem. Photobiol.* **55**, 58S.
- Zaidi, S. I. A., N. L. Oleinick, M. T. Zaim and H. Mukhtar (1993) Apoptosis during photodynamic therapy-induced ablation of RIF-1 tumors in C3H mice: Electron microscopic, histopathologic and biochemical evidence. *Photochem. Photobiol.* **58**, 771-776.
- Zaidi, S. I. A., R. Agarwal, G. Eichler, B. D. Rihter, M. E. Kenney and H. Mukhtar (1993) Photodynamic effects of new silicon phthalocyanines: in vitro studies utilizing rat hepatic microsomes and human erythrocyte ghosts as model membrane sources. *Photochem. Photobiol.* **58**, 204-210.
- Zhang, W.-W. and J. A. Roth (1994) Anti-oncogene and tumor suppressor gene therapy-examples from a lung cancer animal model. *In Vivo* **8**, 755-770.
- Zhang, Z., K. Vuori, J. C. Reed and E. Ruoslahti (1995) The $\alpha 5 \beta 1$ integrin supports survival of cells on fibronectin and up-regulates Bcl-2 expression. *Proc. Natl. Acad. Sci. USA* **92**, 6161-6165.
- Zharkova, N. N., D. N. Kozlev, V. V. Smirnov, V. V. Sokolov, V. I. Chissov, E. V. Filonenko, D. G. Sukhin, M. G. Galpern and G. N. Vorozhtsov (1994) Fluorescence observations of patients in the course of photodynamic therapy of cancer with photosensitizer PHOTOLENS. *Proc. SPIE* **2325**, 400-403.
- Zhou C., C. Milanesi and G. Jori (1988) An ultrastructural comparative evaluation of tumours photosensitized by porphyrins administered in aqueous solution, bound to liposomes or to lipoproteins. *Photochem. Photobiol.* **48**, 487-492.
- Zhou C., C. Shynji, D. Jinsheng, L. Junlin, G. Jori and C. Milanesi (1996) Apoptosis of mouse MS-2 fibrosarcoma cells induced by photodynamic therapy with Zn(II)-phthalocyanine. *J. Photochem. Photobiol. B: Biol.* **33**, 219-223.
- Zickendraht, C. and E. J. Koller (1959) New phthalocyanine dyestuffs. *U.S. Patent 2,897,207* (C.A. 54:P2767e)
- Zuk, M. M., B. D. Rihter, M. E. Kenney, M. A. Rodgers and M. Kreimer-Birnbaum (1996) Effect of delivery system on the pharmacokinetics and tissue distribution of bis(di-isobutyl octadecylsiloxy)silicon 2,3 naphthalocyanine (isoBOSINC), a photosensitizer for tumor therapy. *Photochem. Photobiol.* **63**, 132-140.

Appendix 1.

Role of Activated Oxygen Species in Photodynamic Therapy

Wesley M. Sharman, Cynthia M. Allen and Johan E. van Lier (2000)
Meth in Enzymol., **319**, 376-400.

Pour l'article complet voir la copie papier à la Bibliothèque des sciences de la santé
Section Monographie WB 480 A57 2001

Appendix 2.

Photodynamic Therapeutics: Basic Principles and clinical applications

Wesley M. Sharman, Cynthia M. Allen and Johan E. van Lier (1999)
Drug Discovery Today 4, 507-517.

Pour l'article complet voir la copie papier à la Bibliothèque des sciences de la santé
Section Monographie WB 480 A57 2001

Appendix 3.

Current Status of Phthalocyanines in the Photodynamic Therapy of Cancer

Cynthia M. Allen, Wesley M. Sharman and Johan E. van Lier (2001)
J. Porphyrins Phthalocyanines 5, 161-169.

Pour l'article complet voir la copie papier à la Bibliothèque des sciences de la santé
Section Monographie WB 480 A57 2001